

CYCLE DECOMPOSITIONS IN k -UNIFORM HYPERGRAPHS

ALLAN LO, SIMÓN PIGA, AND NICOLÁS SANHUEZA-MATAMALA

ABSTRACT. We show that k -uniform hypergraphs on n vertices whose codegree is at least $(2/3 + o(1))n$ can be decomposed into tight cycles, subject to the trivial divisibility conditions. As a corollary, we show those graphs contain tight Euler tours as well. In passing, we also investigate decompositions into tight paths.

In addition, we also prove an alternative condition for building absorbers for edge-decompositions of arbitrary k -uniform hypergraphs, which should be of independent interest.

1. INTRODUCTION

Given a k -uniform hypergraph H , a *decomposition of H* is collection of subgraphs of H such that every edge is covered exactly once. If all these subgraphs are isomorphic copies of the same k -uniform graph F , we say H has an F -*decomposition*, and that H is F -*decomposable*. We refer the reader to the survey of Glock, Kühn, and Osthus [8] for a recent account on extremal aspects of hypergraph decomposition problems. Here we investigate hypergraph decompositions into tight cycles.

Given $\ell > k \geq 2$, the k -uniform tight cycle of length ℓ , denoted by $C_\ell^{(k)}$, is the k -graph whose vertices are $\{v_1, \dots, v_\ell\}$ and its edges are $\{v_i, v_{i+1}, \dots, v_{i+k-1}\}$ for all $i \in \{1, \dots, \ell\}$, with the subindices understood modulo ℓ . Given a vertex set $S \subseteq V(H)$, we define the *degree* $\deg_H(S)$ of S as the number of edges of H which contain S . Given a vertex $v \in V(H)$, we define the *degree of v* as the degree of $\{v\}$. Given some $0 \leq i < k$, we let $\delta_i(H)$ (and $\Delta_i(H)$) be the minimum (and maximum, respectively,) value of $\deg_H(S)$ taken over all i -sets of vertices S . We call $\delta_{k-1}(H)$ the *minimum codegree of H* and sometimes we will write just $\delta(H)$ if k is clear from context.

We say that a k -graph H is $C_\ell^{(k)}$ -*divisible* if $|E(H)|$ is divisible by ℓ and the degree of every vertex of H is divisible by k . Clearly, being $C_\ell^{(k)}$ -divisible is a necessary condition to admit an $C_\ell^{(k)}$ -decomposition, but in general it is not a sufficient condition. We are interested in extremal questions of the sort: which conditions on the minimum degree of large $C_\ell^{(k)}$ -divisible graphs ensure the existence of $C_\ell^{(k)}$ -decompositions? Given a k -graph F , we define the F -*decomposition threshold* δ_F be the least $d > 0$ such that for every $\varepsilon > 0$, there exists n_0 such that any F -divisible k -graph H on $n \geq n_0$ vertices with $\delta_{k-1}(H) \geq (d + \varepsilon)n$ admits an F -decomposition.

In this paper, we are interested in $\delta_{C_\ell^{(k)}}$. For $k = 2$, k -graphs are just graphs, tight cycles are just graph cycles, and minimum codegree is just minimum degree, and here much more is known about the values of $\delta_{C_\ell^{(2)}}$. Barber, Kühn, Lo, and Osthus [3] show that $\delta_{C_4^{(2)}} = 2/3$ and for each even $\ell \geq 6$, $\delta_{C_\ell^{(2)}} = 1/2$. Taylor [16] proved exact minimum degree conditions which yield decompositions into cycles of length ℓ in large graphs, for $\ell = 4$ and every even $\ell \geq 8$. For odd values of ℓ , the situation is different. Joos and Kühn [11] showed that $\delta_{C_\ell^{(2)}} = 1/2 + c_\ell$, where c_ℓ is a sequence of non-zero numbers depending on ℓ only, which satisfy $c_\ell \rightarrow 0$ when $\ell \rightarrow \infty$.

For $k = 3$, the last two authors [14] showed that $\delta_{C_\ell^{(3)}} = 2/3$ for sufficiently large ℓ . In fact, they show that the constant ‘2/3’ is also sharp for the more general problem

The research leading to these results was supported by ANID-Chile through the FONDECYT Iniciación N°11220269 grant (N. Sanhueza-Matamala) and EPSRC, grant no. EP/V002279/1 (A. Lo and S. Piga) and EP/V048287/1 (A. Lo). There are no additional data beyond that contained within the main manuscript.

of decomposing hypergraphs into tight cycles of possibly different lengths, which we describe now.

A (*tight*) *cycle-decomposition* of a k -graph H is an edge partition of H into tight cycles (of possibly different lengths). A condition which is easily seen to be necessary to admit a cycle-decomposition is that the degree of every vertex of H is divisible by k . We define the *cycle-decomposition threshold* $\delta_{\text{cycle}}^{(k)}$ be the least $d > 0$ such that for every $\varepsilon > 0$, there exists n_0 such that any k -graph H on $n \geq n_0$ vertices with $\delta_{k-1}(H) \geq (d + \varepsilon)n$ such that every vertex of H is divisible by k admits a cycle-decomposition. Note that $\delta_{\text{cycle}}^{(2)} = 0$ as every graph with even degrees admits a cycle decomposition. For $k = 3$, the last two authors [14] showed that $\delta_{\text{cycle}}^{(3)} = 2/3$. Glock, Kühn and Osthus [8, Conjecture 5.5] posed the following conjecture for $k \geq 3$.

Conjecture 1.1 (Glock, Kühn and Osthus). *For $k \geq 3$, $\delta_{\text{cycle}}^{(k)} \leq (k - 1)/k$.*

Cycle decompositions are also related with generalisations of Euler tours to hypergraphs. An *Euler tour* in a k -graph H is a sequence of (possibly repeating) vertices $v_1 \cdots v_m$ such that each k cyclically consecutive vertices forms an edge of H , and all edges of H appear uniquely in this way. Similarly, we define the *Euler tour threshold* $\delta_{\text{Euler}}^{(k)}$ be the least $d > 0$ such that for every $\varepsilon > 0$, there exists n_0 such that any k -graph H on $n \geq n_0$ vertices with $\delta_{k-1}(H) \geq (d + \varepsilon)n$ such that every vertex of H is divisible by k admits an Euler tour. Chung, Diaconis, and Graham [5] conjectured that every large $K_n^{(k)}$ such that every vertex of H has degree divisible by k admits an Euler tour. Glock, Joos, Kühn, and Osthus [6] confirmed this conjecture and showed the existence of Euler tours in suitable hypergraphs by using results on cycle decompositions, which in particular show $\delta_{\text{Euler}}^{(k)} < 1$ for all k . For $k = 2$, it is easy to see that $\delta_{\text{Euler}}^{(2)} = 1/2$ (as $\delta(H) \geq |V(H)|/2$ is needed to ensure that the graph H is connected), and examples show that $\delta_{\text{Euler}}^{(k)} \geq 1/2$ holds for all $k \geq 3$ [6, Section 1.3]. The following conjecture for all $k \geq 3$ was posed.

Conjecture 1.2 (Glock, Kühn, and Osthus [8]). *For $k \geq 3$, $\delta_{\text{Euler}}^{(k)} \leq (k - 1)/k$.*

It was first conjectured that $\delta_{\text{Euler}}^{(k)} = 1/2$ for all $k \geq 3$ in [6], but this was disproven by the last two authors [14] by showing that $\delta_{\text{Euler}}^{(3)} = 2/3$.

Our main result bounds $\delta_{C_\ell}^{(k)}$ for every $k \geq 2$ and each sufficiently large ℓ .

Theorem 1.3. *For every $k \geq 3$ there exists an $\ell_0 \in \mathbb{N}$ such that for every $\ell \geq \ell_0$ it holds that $\delta_{C_\ell}^{(k)} \leq 2/3$.*

The case when $k = 3$ already appears in [14]. For $k \geq 4$, we do not know if the constant ‘ $2/3$ ’ appearing in Theorem 1.3 is best-possible. We discuss lower bounds in Section 2.

In order to prove Theorem 1.3, we also find the decomposition threshold for tight paths. Given $\ell > k \geq 2$, the *k -uniform tight path on ℓ vertices*, denoted by $P_\ell^{(k)}$, is the k -graph whose vertices are $\{v_1, \dots, v_\ell\}$ and its edges are $\{v_i, v_{i+1}, \dots, v_{i+k-1}\}$ for all $i \in \{1, \dots, \ell - k + 1\}$. A k -graph H is *$P_\ell^{(k)}$ -divisible* if $|E(H)|$ is divisible by $\ell - k + 1$. We prove that $\delta_{P_\ell^{(k)}} = 1/2$.

Theorem 1.4. *For every $k \geq 3$ and $\ell \geq k + 1$, $\delta_{P_\ell^{(k)}} = 1/2$.*

Using the techniques we apply to prove our main result, we can give bounds on $\delta_{\text{cycle}}^{(k)}$ and $\delta_{\text{Euler}}^{(k)}$, which in particular prove Conjectures 1.1 and 1.2 in a strong sense. In fact, we also prove that both thresholds are always equal.

Theorem 1.5. *For all $k \geq 3$, $1/2 \leq \delta_{\text{Euler}}^{(k)} = \delta_{\text{cycle}}^{(k)} \leq \inf_{\ell > k} \{\delta_{C_\ell}^{(k)}\} \leq 2/3$.*

1.1. Proof ideas. Our proof uses the ‘iterative absorption’ framework to tackle decomposition problems in hypergraphs; see [2] for an introduction. The proof of the main result (Theorem 1.3) has three ingredients: an Absorber lemma, a Vortex lemma, and a Cover-down lemma. The Vortex lemma gives a sequence of subsets $V(H) = U_0 \supseteq$

$U_1 \supseteq \dots \supseteq U_t$, and U_t has size independent of $n = |V(H)|$. The Absorber lemma gives a small subgraph $A \subseteq H$ such that for any $C_\ell^{(k)}$ -divisible leftover L , the k -graph $A \cup L$ has a cycle decomposition. This reduces the problem to the search of a cycle packing in $H' = H - A$ which only has uncovered edges in U_t . This is found using the Cover-down lemma: in the i th step we find a collection of edge-disjoint cycles which covers all edges in $H'[U_i] - H'[U_{i+1}]$ but only uses few edges in $H'[U_{i+1}]$, this allows the process to be iterated.

Our proof of the Cover-down lemma requires a result of Joos and Kühn on ‘fractional decompositions’ [11], and a detour which finds and uses (tight) path decompositions.

Most of the work is required to prove the Absorber lemma. We follow the approach of [14], where absorbers are built by first finding ‘tour-trail decompositions’ of the leftover graphs. These decompositions consist on edge-disjoint subgraphs, each of which forms a tour or a trail. It turns out that it is simple to build absorbers if the leftover can be decomposed into tours. The goal is then to modify the leftover via the addition of *gadgets*, these will suitably modify a given tour-trail decomposition in steps, so that at the end no trails remain. We also prove an alternative condition for the existence of absorbers, see Lemma 3.3, which should be of independent interest.

In this high-level description, this is the same outline used to find cycle decompositions when $k = 3$ in [14], but the proof for $k > 3$ requires several non-trivial modifications. This is specially true in the construction of the absorbers, which is way more involved than in the $k = 3$ case, and can be considered the main new contribution of the paper.

1.2. Organisation. In Section 2 we give new lower bounds for the $C_\ell^{(k)}$ -decomposition threshold, for certain values of k and ℓ .

In Section 3, we establish a connection between the notion of *transformers* and *absorbers*. In Section 4 we explain the iterative absorption method, including the statements of their key lemmas. At the end of this section we prove Theorem 1.3.

Sections 5 to 9 are devoted to the proofs of the lemmas used in the iterative absorption. Section 5 contains the proof of the Vortex lemma. The proof of the Absorber lemma is the main technical part of our paper, and its proof spans Sections 6, 7, and 8. The proof of the Cover-down lemma appears in Section 9. We prove Theorem 1.4 in Section 9.1.

In Section 10 we provide the necessary lemmas for the proof of Theorem 1.5. We finish in Section 11 with remarks and questions.

1.3. Notation. Let $[n] = \{1, \dots, n\}$. Since isolated vertices make no difference in our context, we usually do not distinguish from a hypergraph $H = (V(H), E(H))$ and its set of edges $E(H)$. For a subset $U \subseteq V(H)$, we write $H \setminus U$ to mean the subgraph of H obtained by deleting vertices in U . We write $H[U] = H \setminus (V(H) \setminus U)$. For a k -graph G , we write $H - G = (V(H), E(H) \setminus E(G))$. We will suppress brackets and commas to refer to k -tuples of vertices when they are considered as edges of a hypergraph. For instance, for $v_1, \dots, v_k \in V(H)$, $v_1 \dots v_k \in H$ means that the edge $\{v_1, \dots, v_k\}$ is in $E(H)$. For a vertex set $S \subseteq V(H)$, the *neighbourhood* $N_H(S)$ of S is the set of vertex sets $T \subseteq V(H) \setminus S$ such that $S \cup T \in H$. Given $U \subseteq V(H)$, define $N_H(S, U) = N_{H[S \cup U]}(S)$. The degrees $\deg_H(S)$ and $\deg(S, U)$ correspond to $|N_H(S)|$ and $|N_H(S, U)|$, respectively. We suppress H if it can be deduced from context.

We also use the following notation. Given $k \geq 2$ and $r \geq 1$, and a k -graph H , define $\delta^{(r)}(H)$ to be the minimum of $|N(e_1) \cap N(e_2) \cap \dots \cap N(e_r)|$ among all possible choices of r different $(k-1)$ -sets of vertices e_1, \dots, e_r . More generally, given a set of vertices $U \subseteq V(H)$, we also define $\delta^{(r)}(H, U)$ as the minimum of $|U \cap N(e_1) \cap N(e_2) \cap \dots \cap N(e_r)|$ among all possible choices of r different $(k-1)$ -sets of vertices e_1, \dots, e_r .

We will use hierarchies in our statements. The phrase “ $a \ll b$ ” means “for every $b > 0$, there exists $a_0 > 0$, such that for all $0 < a \leq a_0$ the following statements hold”. We

implicitly assume all constants in such hierarchies are positive, and if $1/a$ appears we assume a is an integer.

Suppose that Lemma A states that a k -graph H contains a subgraph J . We write ‘apply Lemma A and obtain edge-disjoint subgraphs J_1, \dots, J_ℓ ’ to mean that ‘for each $i \in [\ell]$, we apply Lemma A to $H - \bigcup_{j \in [i-1]} J_j$ to obtain J_i ’. Note that $H - \bigcup_{j \in [i-1]} J_j$ will also satisfy the condition of Lemma A, but we will not check them explicitly. Furthermore, suppose that we have already found a subgraph H' of H and we say that ‘apply Lemma A and obtain subgraph J such that $V(J) \setminus U$ are new vertices’ to mean the ‘we apply Lemma A to $H - (V(H') \setminus U)$ to obtain J ’.

2. LOWER BOUNDS

Given a k -graph H , let $\mathcal{C}_\ell(H)$ be the family of all $C_\ell^{(k)}$ in H and $\mathcal{C}_\ell(H, e)$ the family of ℓ -cycles containing a fixed edge $e \in H$. A *fractional $C_\ell^{(k)}$ -decomposition* of H is a function $\omega : \mathcal{C}_\ell(H) \rightarrow [0, 1]$ such that, for every edge $e \in H$, $\sum_{C \in \mathcal{C}_\ell(H, e)} \omega(C) = 1$. We define the *fractional $C_\ell^{(k)}$ -decomposition threshold* $\delta_{C_\ell^{(k)}}^*$ be the least $d > 0$ such that, for every $\varepsilon > 0$, there exists n_0 such that any k -graph H on $n \geq n_0$ vertices with $\delta_{k-1}(H) \geq (d + \varepsilon)n$ admits a fractional $C_\ell^{(k)}$ -decomposition.

Here, we give lower bounds on the parameter $\delta_{C_\ell^{(k)}}^*$. Joos and Kühn [11] showed that $\delta_{C_\ell^{(k)}}^* \geq \frac{1}{2} + \frac{1}{(k-1+2/k)(\ell-1)}$ holds for each $k \geq 2$ and ℓ not divisible by k . We give new bounds, which remove the dependency on k .

Proposition 2.1. *Let $1 \leq i < k < \ell$ with ℓ not divisible by k and $r = k/\gcd(k, \ell)$. Let $I_{\text{free}} = \{0 \leq i \leq k : i \not\equiv 0 \pmod{r}\}$, $I_{\text{odd}} = \{0 \leq i \leq k : i \not\equiv 0 \pmod{2}\}$ and $I_{\text{even}} = \{0 \leq i \leq k : i \equiv 0 \pmod{2}\}$. Then*

$$(2.1) \quad \delta_{C_\ell^{(k)}}^* \geq \frac{1}{2} + \frac{1}{2^k(\ell-1)} \max \left\{ \sum_{i \in I_{\text{free}} \cap I_{\text{odd}}} \binom{k}{i}, \sum_{i \in I_{\text{free}} \cap I_{\text{even}}} \binom{k}{i} \right\} \geq \frac{1}{2} + \frac{1}{4(\ell-1)}.$$

Our constructions are based on [9, Proposition 3.1]. Given vertex-disjoint sets A, B and $0 \leq i \leq k$, we let $H_i^{(k)}(A, B)$ be the k -graph on $A \cup B$ such that $e \in H_i$ if and only if $|e \cap B| = i$. We need the following observation.

Proposition 2.2. *Let $1 \leq i < k < \ell$ and $d = \gcd(k, \ell)$. Let A and B be disjoint vertex sets, and let $H_i = H_i^{(k)}(A, B)$. Then H_i is $C_\ell^{(k)}$ -free for all $k - i \not\equiv 0 \pmod{k/d}$.*

Proof. Suppose $\ell > k$ is such that $v_1 \cdots v_\ell$ are the vertices of a copy of $C_\ell^{(k)}$ in H_i . Letting $d = \gcd(k, \ell)$, we shall show that k/d divides $k - i$. For all $j \in [\ell]$, let $\phi_j \in \{A, B\}$ be such that $v_j \in \phi_j$ and let $\phi_{\ell+j} = \phi_j$. Note that $\phi_j = \phi_{j+k}$ for all $j \in [\ell]$. Hence, $\phi_{j+d} = \phi_j$ for all $j \in [\ell]$. Thus

$$k - i = |\{v_1, \dots, v_k\} \cap A| = |\{j \in [k] : \phi_j = A\}| \in \{k/d, 2k/d, \dots, k\}$$

as required. \blacksquare

We say a k -graph H on n vertices admits an η -approximate F -decomposition if it has a collection of edge-disjoint copies of F covering all but ηn^k edges. By a result of Rödl, Schacht, Siggers, and Tokushige [15], any bound on the codegree of k -graphs not containing η -approximate decompositions, for arbitrary small η , is essentially equivalent to bound the corresponding numbers for fractional $C_\ell^{(k)}$ -decomposition. Thus, we will focus on the former.

Proof of Proposition 2.1. Let n be sufficiently large. Let A and B be disjoint vertex sets each of size $\lfloor n/2 \rfloor$ and $\lceil n/2 \rceil$ respectively. For each $0 \leq i \leq k$, let $H_i = H_i^{(k)}(A, B)$. Note that

$$(2.2) \quad \frac{|H_i|}{\binom{n}{k}} = \frac{1}{2^k} \binom{k}{i} + o(1).$$

Let $d = \gcd(k, \ell)$. Since ℓ is not divisible by k , then $r = k/d > 1$. It holds that $k-i \not\equiv 0 \pmod r$ if and only if $i \not\equiv 0 \pmod r$. Let $H_{\text{odd}} = \bigcup_{i \in I_{\text{odd}}} H_i$ and $H_{\text{even}} = \bigcup_{i \in I_{\text{even}}} H_i$. Note that $\delta(H_{\text{odd}}), \delta(H_{\text{even}}) \geq n/2 - k$. From (2.2) it follows that both $|H_{\text{odd}}|$ and $|H_{\text{even}}|$ have size $(\frac{1}{2} + o(1))\binom{n}{k}$.

Let $H'_{\text{odd}} = \bigcup_{i \in I_{\text{odd}} \cap I_{\text{free}}} H_i$ and $H'_{\text{even}} = \bigcup_{i \in I_{\text{even}} \cap I_{\text{free}}} H_i$. Note that by (2.2), we have

$$(2.3) \quad \frac{|H'_{\text{odd}}|}{\binom{n}{k}} = \frac{1}{2^k} \sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i} + o(1) \quad \text{and} \quad \frac{|H'_{\text{even}}|}{\binom{n}{k}} = \frac{1}{2^k} \sum_{i \in I_{\text{even}} \cap I_{\text{free}}} \binom{k}{i} + o(1).$$

Observe that given odd numbers $i \neq j$ there no tight path in H_{odd} connecting an edge from H_i with an edge of H_j . Therefore, Proposition 2.2 yields that no edge in H'_{odd} is contained in a copy of $C_\ell^{(k)}$ in H_{odd} . Let

$$p = \frac{2|H'_{\text{odd}}|}{(\ell-1)\binom{n}{k}},$$

and suppose $\eta > 0$ is given. Consider H_{even}^* to be a sub- k -graph of H_{even} such that $\delta(H_{\text{even}}^*) \geq (p - 4\eta)n/2$ and

$$(2.4) \quad |H_{\text{even}}^*| \leq (p - 2.1\eta)|H_{\text{even}}| < |H'_{\text{odd}}|/(\ell-1) - \eta n^k.$$

Such a sub- k -graph can be obtained by taking random edges from H_{even} independently with probability $p - 3\eta$.

We claim $H = H_{\text{odd}} \cup H_{\text{even}}^*$ does not admit an η -approximate $C_\ell^{(k)}$ -decomposition. Since no edge of H'_{odd} is contained in copy of $C_\ell^{(k)}$ in H_{odd} , each $C_\ell^{(k)}$ containing an edge in H'_{odd} must contain at least one edge in H_{even}^* . Therefore, if H contains an η -approximate $C_\ell^{(k)}$ -decomposition, then we have

$$(\ell-1)|H_{\text{even}}^*| \geq |H'_{\text{odd}}| - \eta|H| \geq |H'_{\text{odd}}| - \eta n^k,$$

contradicting (2.4). Note that $\delta(H) \geq \delta(H_{\text{odd}}) + \delta(H_{\text{even}}^*) \geq (1 + p - 4.5\eta)n/2$. Therefore, from (2.3), letting n tend to infinity, and η tend to zero, we deduce that

$$\delta_{C_\ell^{(k)}}^* \geq \lim_{n \rightarrow \infty} \frac{1}{2}(1 + p) = \frac{1}{2} + \frac{1}{2^k(\ell-1)} \sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i}.$$

An analogous construction, selecting $H_{\text{odd}}^* \subseteq H_{\text{odd}}$ as a random set of the appropriate size with respect to H'_{even} , gives that

$$\delta_{C_\ell^{(k)}}^* \geq \frac{1}{2} + \frac{1}{2^k(\ell-1)} \sum_{i \in I_{\text{even}} \cap I_{\text{free}}} \binom{k}{i},$$

and therefore we have

$$\delta_{C_\ell^{(k)}}^* \geq \frac{1}{2} + \frac{1}{2^k(\ell-1)} \max \left\{ \sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i}, \sum_{i \in I_{\text{even}} \cap I_{\text{free}}} \binom{k}{i} \right\}.$$

which gives the first inequality of (2.1).

To bound this last term, note that

$$\sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i} + \sum_{i \in I_{\text{even}} \cap I_{\text{free}}} \binom{k}{i} = \sum_{1 \leq i \leq k, i \not\equiv 0 \pmod r} \binom{k}{i} \geq 2^{k-1}.$$

The last inequality follows since $\sum_{i \in [k]: i \equiv 0 \pmod r} \binom{k}{i}$ counts the number of sets of $[k]$ of size divisible by r , and we recall that $r > 1$. If $\mathcal{P}_r \subseteq \mathcal{P}([k])$ is that family, then $X \mapsto X \Delta \{1\}$ is an injection from \mathcal{P}_r to $\mathcal{P}([k]) \setminus \mathcal{P}_r$, and thus $|\mathcal{P}_r| \leq |\mathcal{P}([k])|/2 = 2^{k-1}$.

We deduce that $\max \left\{ \sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i}, \sum_{i \in I_{\text{even}} \cap I_{\text{free}}} \binom{k}{i} \right\} \geq 2^{k-2}$, which then yields

$$\delta_{C_\ell^{(k)}}^* \geq \frac{1}{2} + \frac{1}{4(\ell-1)},$$

as desired. \blacksquare

We can get better bounds for some choices of k and ℓ by looking at (2.1) in detail.

Corollary 2.3. *Let $3 \leq k < \ell$ with $\ell \not\equiv 0 \pmod k$. Then*

$$\delta_{C_\ell}^* \geq \begin{cases} \frac{1}{2} + \frac{1}{2(\ell-1)} & \text{if } k/\gcd(\ell, k) \text{ is even,} \\ \frac{1}{2} + \frac{1-2^{-k}}{2(\ell-1)} & \text{if } \gcd(\ell, k) = 1 \text{ and } k \text{ is odd.} \end{cases}$$

Proof. Let $d = \gcd(k, \ell)$ and $k = dr$. If r is even, then $I_{\text{odd}} \cap I_{\text{free}} = I_{\text{odd}}$. Therefore $\sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i} = \sum_{i \in I_{\text{odd}}} \binom{k}{i} = 2^{k-1}$, so $\delta_{C_\ell}^* \geq \frac{1}{2} + \frac{1}{2(\ell-1)}$ follows from Proposition 2.1.

If $d = 1$, then $r = k$, and therefore $I_{\text{free}} = [k-1]$. This implies $\sum_{i \in I_{\text{free}}} \binom{k}{i} = 2^k - 2$ and therefore $\max \left\{ \sum_{i \in I_{\text{odd}} \cap I_{\text{free}}} \binom{k}{i}, \sum_{i \in I_{\text{even}} \cap I_{\text{free}}} \binom{k}{i} \right\} \geq 2^{k-1} - 1$, then $\delta_{C_\ell}^* \geq \frac{1}{2} + \frac{1-2^{-k}}{2(\ell-1)}$ again follows from Proposition 2.1. \blacksquare

Finally, we can get bounds for the non-fractional thresholds $\delta_{C_\ell}^{(k)}$ by modifying the k -graphs we construct in the proof of Proposition 2.1 in such a way that they also are $C_\ell^{(k)}$ -divisible. By removing at most $\ell - 1$ edges it is easy to make the total number of edges divisible by ℓ , so the only real challenge is to make every degree divisible by k . We prove later (Corollary 9.11) that, for each $\varepsilon > 0$ (assuming n sufficiently large), we can find $F \subseteq H$ whose number of edges is divisible by ℓ , $\delta_{k-1}(H - F) \geq \delta_{k-1}(H) - \varepsilon n$, and for each $v \in V(H)$, $\deg_{H-F}(v) \equiv 0 \pmod k$. Thus the graphs we construct in Proposition 2.1 can be modified to be $C_\ell^{(k)}$ -divisible, which implies the following bounds:

Corollary 2.4. *For all $3 \leq k < \ell$ and ℓ not divisible by k ,*

- (i) $\delta_{C_\ell}^{(k)} \geq \frac{1}{2} + \frac{1}{4(\ell-1)}$,
- (ii) *if $k/\gcd(\ell, k)$ is even, then $\delta_{C_\ell}^{(k)} \geq \frac{1}{2} + \frac{1}{2(\ell-1)}$, and*
- (iii) *if $k/\gcd(\ell, k) = 1$ and ℓ is odd, then $\delta_{C_\ell}^{(k)} \geq \frac{1}{2} + \frac{1-2^{-k}}{2(\ell-1)}$.*

3. ABSORBERS VERSUS TRANSFORMERS

In this section, we introduce absorbers and transformers, which are essential tools in the iterative absorption technique. We prove that the existence of absorbers is essentially equivalent to the existence of transformers, and we work with the latter concept in the rest of the paper. We state our results in a general fashion, that is, for F -decompositions into general hypergraphs, not just cycles.

Let F and G be k -graphs. We say that a k -graph A is an F -absorber for G if both A and $A \cup G$ have F -decompositions and $A[V(G)] = \emptyset$. Note that if there is an F -absorber for G , then G is F -divisible. The following definition describe k -graphs containing absorbers in a robust way.

Definition 3.1. Let $\eta : \mathbb{N} \rightarrow \mathbb{N}$ be a function. We say that a k -graph H on n vertices is (F, m_G, m_W, η) -absorbing if, for all F -divisible subgraphs G of H with $|V(G)| \leq m_G$ and $W \subseteq V(H) \setminus V(G)$ with $|W| \leq m_W - \eta(|V(G)|)$, $H \setminus W$ contains an F -absorber A for G with $|A| \leq \eta(|V(G)|)$.

We will use so-called transformers to construct absorbers. The rôle of transformers allows us to replace G with a ‘homomorphic copy’ G' of G . Given k -graphs G and G' , a function $\phi : V(G) \rightarrow V(G')$ is an *edge-bijective homomorphism from G to G'* if we have $G' = \{\phi(v_1) \dots \phi(v_k) : v_1 \dots v_k \in G\}$. A $(G, G'; F)$ -transformer is a k -graph T such that $T \cup G$ and $T \cup G'$ are F -decomposable and $T[V(G)] \cup T[V(G')]$ is empty. The following definition is analogous to Definition 3.1 but for transformers.

Definition 3.2. Let $\eta : \mathbb{N} \rightarrow \mathbb{N}$ be an increasing function with $\eta(x) \geq x$. We say that a k -graph H on n vertices is (F, m_G, m_W, η) -transformable if, for vertex-disjoint homomorphic F -divisible subgraphs G, G' of H and $W \subseteq V(H) \setminus V(G \cup G')$

and $|V(G)|, |V(G')| \leq m_G$ and $|W| \leq m_W - \eta(|V(G)|)$, $H \setminus W$ contains a $(G, G'; F)$ -transformer T with $|T| \leq \eta(\max\{|V(G)|, |V(G')|\})$.

It is not difficult to see that if a k -graph H is (F, m_G, m_W, η) -absorbing, then H is also $(F, m_G, m_W, 2\eta)$ -transformable (see proof of Lemma 3.3). In fact, the converse is true with different constants as long as there are enough copies of F in H .

Lemma 3.3. *Let $\eta : \mathbb{N} \rightarrow \mathbb{N}$ be an increasing function with $\eta(x) \geq x$ and $m_W, m_G \geq 0$. Let F be a k -graph and $u_1 \cdots u_k \in F$. Let H be a k -graph such that, for any distinct $v_1, \dots, v_k \in V(H)$ and $W \subseteq V(H) \setminus \{v_i : i \in [k]\}$ with $|W| \leq m_W$, $(H \cup \{v_1, \dots, v_k\}) \setminus W$ contains a copy of F with u_i mapped to v_i for all $i \in [k]$. Then if H is (F, m_G, m_W, η) -absorbing then H is $(F, m_G, m_W, 2\eta)$ -transformable. Moreover, if H is (F, m_G, m_W, η) -transformable, then H is (F, m_G, m_W, η') -absorbing for some increasing function $\eta' : \mathbb{N} \rightarrow \mathbb{N}$ with $\eta'(x) \geq x$.*

For k -graphs G and H and $q \in \mathbb{N}$, we write $G + qH$ to be the vertex-disjoint union of G and q copies of H . We now show that, by adding q vertex-disjoint copies of F to G , the k -graph $G + qF$ has an edge-bijective homomorphism to $K_m^{(k)}$. We will require the following theorem regarding the existence of F -decompositions in high codegree k -graphs. Recall that δ_F is the F -decomposition threshold.

Theorem 3.4 (Glock, Kühn, Lo and Osthus [7]). *For all k -graphs F , there exists a constant $c_F > 0$ such that $\delta_F \leq 1 - c_F$.*

A subsequent alternative proof of Theorem 3.4 was given by Keevash [12].

Lemma 3.5. *Let F be a k -graph. Then, for all $t \in \mathbb{N}$, there exist integers $q = q(t)$ and $m = m(t)$ such that, for any F -divisible k -graph G with $|G| = t$, there exists an edge-bijective homomorphism from $G + qF$ to $K_m^{(k)}$.*

Proof. Let $1/m \ll 1/t, 1/k$ be such that $K_m^{(k)}$ is F -divisible. Let G' be an isomorphic copy of G with $V(G') \subseteq V(K_m^{(k)})$. Clearly there is an edge-bijective homomorphism from G to G' . To prove the lemma, it suffices to show that $H = K_m - G$ is F -decomposable and set $q = |H|/|F|$. Both G and $K_m^{(k)}$ are F -divisible, so is H . Note that $|V(G)| \leq k|G| \leq kt$, so $\delta(H) \geq (1 - kt/m)m$. By Theorem 3.4, H has an F -decomposition. ■

We now sketch the how to construct absorber from transformers, that is, the backwards direction of the proof of Lemma 3.3. Let G be an F -divisible k -graph with t edges. By Lemma 3.5, $G + qF$ has an edge-bijective homomorphism to $K_m^{(k)}$. Hence there exists a $(G + qF, K_m^{(k)}; F)$ -transformer T . Note that $T_1 = T \cup qF$ is a $(G, K_m^{(k)}; F)$ -transformer. Since m and q only depend on $t = |G|$, by replacing G with $s = t/|F|$ copies of F , we can find a $(sF, K_m^{(k)}; F)$ -transformer T_2 . Then $T_1 \cup K_m^{(k)} \cup T_2$ is a $(G, sF; F)$ -transformer and so $T_1 \cup K_m^{(k)} \cup T_2$ is an F -absorber for G .

However, an obvious obstacle with this approach is that H may not contain any such large clique $K_m^{(k)}$. To overcome this problem, we consider the extension operator ∇ (which was introduced in [7, Definition 8.13]). Fix an edge $u_1 \cdots u_k \in F$. Consider any distinct vertices $v_1, \dots, v_k \in V(H)$. Define $\nabla_{F, u_1 \dots u_k}(v_1 \dots v_k)$ to be a copy of $F - u_1 \cdots u_k$ with v_i playing the rôles of u_i . For a k -graph G on $V(G) \subseteq V(H)$, define $\nabla_{F, u_1 \dots u_k}(G)$ to be the union of $\bigcup_{e \in G} \nabla_{F, u_1 \dots u_k}(e)$, where the ordering of each edge $e \in G$ will be clear from the context and $V(\nabla_{F, u_1 \dots u_k}(e)) \setminus e$ are new vertices. Note that $G \cup \nabla_{F, u_1 \dots u_k}(G)$ is F -decomposable. The hypothesis of Lemma 3.3 implies that $\nabla_{F, u_1 \dots u_k}(G)$, $\nabla_{F, u_1 \dots u_k}(K_m^{(k)})$ and $\nabla_{F, u_1 \dots u_k}(sF)$ exist. We then construct transformers between them to obtain an F -absorber for G .

Proof of Lemma 3.3. First suppose that H is (F, m_G, m_W, η) -absorbing. Let G and G' be vertex-disjoint F -divisible subgraphs of H with $|V(G)|, |V(G')| \leq m_G$. Let $W \subseteq V(H) \setminus V(G \cup G')$ with $|W| \leq m_W - 2\eta(\max\{|V(G)|, |V(G')|\})$. By the property of being (F, m_G, m_W, η) -absorbing, $H \setminus (W \cup V(G'))$ contains an F -absorber A_1 for G

with $A_1[V(G)] = \emptyset$ and $|V(A_1)| \leq \eta(|V(G')|)$. Also, $H \setminus (W \cup V(A_1))$ contains an F -absorber A_2 for G' with $A_2[V(G')] = \emptyset$ and $|A_2| \leq \eta(|V(G')|)$. Let $T = A_1 \cup A_2$. Note that $T \cup G$ and $T \cup G'$ have F -decompositions and $T[V(G \cup G')] = \emptyset$. Hence T is an $(G_1, G_2; F)$ -transformer. Moreover,

$$|V(T)| = |V(A_1)| + |V(A_2)| \leq \eta(|V(G)|) + \eta(|V(G')|) \leq 2\eta(\max\{|V(G)|, |V(G')|\}).$$

So H is $(F, m_G, m_W, 2\eta)$ -transformable.

Now suppose that H is (F, m_G, m_W, η) -transformable. Let $q(t)$ and $m(t)$ be the functions given by Lemma 3.5. Let η' be the function given by

$$\eta'(x) = 2\eta \left(|V(F)| \binom{m(x)}{k} \right).$$

We now show that H is (F, m_G, m_W, η') -absorbing.

Let G be an F -divisible subgraph of H with $|V(G)| \leq m_G$ and $W \subseteq V(H) \setminus V(G)$ with $|W| \leq m_W - \eta'(m_G)$. Let

$$q_1 = q(|G|), \quad m_1 = m(|G|), \quad q_2 = \binom{m_1}{k} / |F|.$$

Let $(q_1 + q_2)F$ be in $H \setminus (V(G) \cup W)$, which exists by our assumption of H . Let $G_1 = G + q_1F$ and $G_3 = q_2F$. Hence, G_1 and G_3 are vertex-disjoint and are in $H \setminus W$. Let $V' = \{v'_1, \dots, v'_{m_1}\} \subseteq V(H) \setminus (V(G_1 \cup G_3) \cup W)$. Consider a $K_{m_1}^{(k)}$ on V' , which may not exist in H . By Lemma 3.5, there exists an edge-bijective homomorphism ϕ_j from G_j to $K_m^{(k)}$ for $j \in \{1, 3\}$. Order edges in $K_m^{(k)}$ into $v'_{i_1} \dots v'_{i_k}$ such that $i_1 < \dots < i_k$. By ϕ_1 and ϕ_3 , this implies an ordering on all edges of $G_1 \cup G_3$. Fix an edge $u_1 \dots u_k \in F$. Let

$$G'_1 = \nabla_{F, u_1 \dots u_k}(G_1), \quad G'_2 = \nabla_{F, u_1 \dots u_k}(K_{m_1}^{(k)}), \quad G'_3 = \nabla_{F, u_1 \dots u_k}(G_3).$$

Let $\ell = |V(F)| \binom{m_1}{k}$. Note that

$$|V(G'_j)| \leq |V(F)| |G'_j| = |V(F)| \binom{m_1}{k} = \ell.$$

By the property of H , H contains vertex-disjoint G'_1, G'_2, G'_3 such that $G'_j[V(G_j)] = \emptyset$. Since H is (F, m_G, m_W, η) -transformable, $H \setminus (W \cup V(G'_3))$ contains a $(G'_1, G'_2; F)$ -transformer T_1 with $|V(T_1)| \leq \eta(\ell)$. Similarly, $H \setminus (W \cup V(G'_1) \cup (T_1 \setminus V(G'_2)))$ contains a $(G'_2, G'_3; F)$ -transformer T_2 with $|V(T_2)| \leq \eta(\ell)$.

Let $A = (G_1 - G) \cup G'_1 \cup T_1 \cup G'_2 \cup T_2 \cup G'_3 \cup G_3$. Recall that $(G_1 - G)$, $G_1 \cup G'_1$, $G'_3 \cup G_3$ and G_3 have F -decompositions. Hence

$$\begin{aligned} A \cup G &= (G_1 \cup G'_1) \cup (T_1 \cup G'_2) \cup (T_2 \cup G'_3) \cup G_3 \text{ and} \\ A &= (G_1 - G) \cup (G'_1 \cup T_1) \cup (G'_2 \cup T_2) \cup (G'_3 \cup G_3) \end{aligned}$$

are F -decomposable. Therefore A is an F -absorber for G . Note that $A[V(G)] = T_1[V(G)] \subseteq T_1[V(G'_1)] = \emptyset$ and

$$|V(A)| \leq |V(T_1)| + |V(T_2)| \leq 2\eta(\ell) = \eta'(|V(G)|).$$

Hence H is (F, m_G, m_W, η') -absorbing. ■

4. ITERATIVE ABSORPTION AND PROOF OF THE MAIN RESULT

The method of iterative absorption is based on three main lemmas: the Vortex lemma, the Absorber lemma, and the Cover-down lemma. We state these lemmas while explaining the general strategy, then we will use them to prove Theorem 1.3. We proof of these lemmas are in Sections 5-9 (Sections 6-8 are dedicated to the Absorber lemma).

A sequence of nested subsets $U_0 \supseteq \dots \supseteq U_t$ of vertices of a k -graph H is a (δ, ξ, m) -vortex for H if

- (V1) $U_0 = V(H)$,
- (V2) for each $i \in [t]$, $|U_i| = \lfloor \xi |U_{i-1}| \rfloor$,
- (V3) $|U_t| = m$,
- (V4) $\delta^{(2)}(H[U_i]) \geq \delta |U_i|$, for each $0 \leq i \leq t$ and
- (V5) $\delta^{(2)}(H[U_i], U_{i+1}) \geq \delta |U_{i+1}|$, for each $0 \leq i < t$.

The Vortex lemma gives us the existence of vortices with the right parameters.

Lemma 4.1 (Vortex lemma). *Let $\delta > 0$ and $1/m' \ll \xi, 1/k$. Let H be a k -graph on $n \geq m'$ vertices with $\delta^{(2)}(H) \geq \delta$. Then H has a $(\delta - \xi, \xi, m)$ -vortex, for some $\lfloor \xi m' \rfloor \leq m \leq m'$.*

Using the properties of such a vortex, we will iteratively find $C_\ell^{(k)}$ -packings covering the edges from $H[U_i]$ in every step, without taking too many edges from the following sets U_{i+1}, \dots, U_t in the vortex. The Cover-down lemma will provide the existence of those packings in every step.

Lemma 4.2 (Cover-down lemma). *For every $k \geq 3$ and every $\alpha > 0$, there is an $\ell_0 \in \mathbb{N}$ such that for every $\mu > 0$ and every $n, \ell \in \mathbb{N}$ with $\ell \geq \ell_0$ and $1/n \ll \mu, \alpha$ the following holds. Let H be a k -graph on n vertices, and $U \subseteq V(H)$ with $|U| = \lfloor \alpha n \rfloor$, and they satisfy*

- (CD₁) $\delta^{(2)}(H) \geq 2\alpha n$,
- (CD₂) $\delta^{(2)}(H, U) \geq \alpha |U|$ and
- (CD₃) $\deg_H(x)$ is divisible by k for each $x \in V(H) \setminus U$.

Then H contains a $C_\ell^{(k)}$ -decomposable subgraph $F \subseteq H$ such that $H - H[U] \subseteq F$ and $\Delta_{k-1}(F[U]) \leq \mu n$.

Finally, after repeated applications of the Cover-down lemma, we only need to consider the edges remaining in $H[U_t]$. For these last edges, we apply the Absorber lemma. This lemma says that the k -graph H is $(C_\ell^{(k)}, m, m, \eta')$ -absorbing, and therefore, it contains an absorber for *any* possible $C_\ell^{(k)}$ -divisible k -graph left as a remainder in U_t (which is of size m).

Lemma 4.3 (Absorber lemma). *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k, 1/m$ with $k \geq 3$ and $\ell \geq 2(k^2 - k) + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq 2\varepsilon n$. Then H is $(C_\ell^{(k)}, m, m, \eta')$ -absorbing for some increasing function $\eta' : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\eta'(x) \geq x$ and independent of ε and n .*

Now we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. We divide the proof into three steps: setting the vortex and absorber, covering down, and using the absorber to conclude. We suppose $\varepsilon, \ell, m', n_0$ are chosen according to the following hierarchy: $1/n_0 \ll 1/m' \ll \varepsilon, 1/\ell \ll 1/k$.

Let H be a $C_\ell^{(k)}$ -divisible k -graph on $n \geq n_0$ with $\delta_{k-1}(H) \geq (2/3 + 8\varepsilon)n$ and observe that this immediately implies that $\delta^{(3)}(H) \geq 8\varepsilon n$ and that $\delta^{(2)}(H) \geq (1/3 + 8\varepsilon)n$. To prove the lemma, it is enough to show that H has a $C_\ell^{(k)}$ -decomposition.

Step 1: Setting the vortex and absorber. By Lemma 4.1, we obtain a $(1/3 - 7\varepsilon, \varepsilon, m)$ -vortex $U_0 \supseteq \dots \supseteq U_t$ in H , for some m satisfying $\lfloor \varepsilon m' \rfloor \leq m \leq m'$.

Let \mathcal{L} be the set of all $C_\ell^{(k)}$ -divisible subgraphs of $H[U_t]$. Clearly, $|\mathcal{L}| \leq 2^{\binom{|U_t|}{k}} \leq 2^{m^k}$. Let $L \in \mathcal{L}$ be arbitrary. Clearly, $\delta^{(3)}(H - H[U_1]) \geq 7\varepsilon n$. By Lemma 4.3 and the choice of constants, we deduce $H - H[U_1]$ is $(C_\ell^{(k)}, m, m, \eta')$ -absorbing for some increasing function $\eta' : \mathbb{N} \rightarrow \mathbb{N}$ which satisfies $\eta'(x) \geq x$. Thus, $H - H[U_1]$ contains an $C_\ell^{(k)}$ -absorber A_L in H' , with $A_L[U_t] = \emptyset$ and $|A_L| \leq \eta'(m)$. We iterate this argument, finding edge-disjoint absorbers $A_{L'} \subseteq H - H[U_1]$, one for each $L' \in \mathcal{L}$. This indeed can be done, since in each step, by removing all the absorbers found so far, we remove at most $|\mathcal{L}|\eta'(m) \leq \varepsilon n/3$ edges overall. Thus $H - H[U_1]$, after removing the already

found absorbers, still satisfies $\delta^{(3)}(H - H[U_1]) \geq 6\epsilon n$ and thus Lemma 4.3 can still be invoked.

Let $A = \bigcup_{L \in \mathcal{L}} A_L \subseteq H - H[U_1]$ be the edge-disjoint union of all absorbers. As argued before, A contains at most $\epsilon n/3$ edges in total. By construction, A is $C_\ell^{(k)}$ -decomposable, and for each $L \in \mathcal{L}$, $A \cup L$ is $C_\ell^{(k)}$ -decomposable. Let $H' = H - A$. Note that $\delta_{k-1}(H') \geq (2/3 + 6\epsilon)n$ and $U_0 \supseteq \dots \supseteq U_t$ is a $(1/3 + 5\epsilon, \epsilon, m)$ -vortex for H' (this is because we have ensured $A \subseteq H - H[U_1]$). Since H and A are $C_\ell^{(k)}$ -divisible, H' is also $C_\ell^{(k)}$ -divisible.

Step 2: Covering down. Now we want to find a $C_\ell^{(k)}$ -packing in H' which covers all edges in $H' - H[U_t]$. For this, we proceed as follows. Let $U_{t+1} = \emptyset$. For each $0 \leq i \leq \ell$ we will find $H_i \subseteq H'[U_i]$ such that

- (a_i) $H' - H_i$ has a $C_\ell^{(k)}$ -decomposition,
- (b_i) $\delta^{(2)}(H_i) \geq (1/3 + 3\epsilon)|U_i|$,
- (c_i) $\delta^{(2)}(H_i, U_{i+1}) \geq (1/3 + 3\epsilon)|U_{i+1}|$, and
- (d_i) $H_i[U_{i+1}] = H'[U_{i+1}]$.

For $i = 0$ this is done by setting $H_0 = H'$. Now, suppose that for $0 \leq i < \ell$ we have found $H_i \subseteq H'[U_i]$ satisfying (a_i)–(d_i), we construct $H_{i+1} \subseteq H'[U_{i+1}]$ satisfying (a_{i+1})–(d_{i+1}). By (a_i), H_i is $C_\ell^{(k)}$ -divisible. Let $H'_i = H_i - H_i[U_{i+1}]$. By (b_i)–(c_i) and $|U_{i+2}| \leq \epsilon|U_{i+1}| \leq \epsilon^2|U_i|$, we have

- (C1) $\delta^{(2)}(H'_i) \geq (1/3 + 2\epsilon)|U_i|$,
- (C2) $\delta^{(2)}(H'_i, U_{i+1}) \geq (1/3 + 2\epsilon)|U_{i+1}|$, and
- (C3) $\deg_{H'_i}(x)$ is divisible by k for each $x \in U_i \setminus U_{i+1}$.

Now we apply Lemma 4.2 with $1/3$, ℓ , ϵ^6 , $|U_i|$, H'_i and U_{i+1} playing the rôles of the parameters α , ℓ , μ , n , H and U . By doing so, we obtain a $C_\ell^{(k)}$ -decomposable subgraph $F_i \subseteq H'_i$ such that $H'_i - H'_i[U_{i+1}] \subseteq F_i$ and $\Delta_{k-1}(F_i[U_{i+1}]) \leq \epsilon^6|U_i|$.

We let $H_{i+1} = H[U_{i+1}] - F_i$, and we now show that it satisfies the required properties. Since $H' - H_{i+1}$ is the edge-disjoint union of $H' - H_i$ and F_i and both are $C_\ell^{(k)}$ -decomposable, we deduce that (a_{i+1}) holds. Note that we have $\Delta_{k-1}(F_i[U_{i+1}]) \leq \epsilon^6|U_i| \leq \epsilon^4|U_{i+1}| \leq \epsilon|U_{i+1}|$. From the definition of $(1/3 + 5\epsilon, \epsilon, m)$ -vortex for H' , we deduce that $\delta^{(2)}(H'[U_{i+1}]) \geq (1/3 + 5\epsilon)|U_{i+1}|$ and $\delta^{(2)}(H'[U_{i+1}], U_{i+2}) \geq (1/3 + 5\epsilon)|U_{i+2}|$. Using this, we are able to deduce that $\delta^{(2)}(H'_{i+1}) \geq \delta^{(2)}(H'[U_{i+1}]) - 2\Delta_{k-1}(F_i[U_{i+1}]) \geq (1/3 + 5\epsilon - 2\epsilon)|U_{i+1}| \geq (1/3 + 3\epsilon)|U_{i+1}|$, and similarly we have $\delta^{(2)}(H'_{i+1}, U_{i+2}) \geq (1/3 + 3\epsilon)|U_{i+2}|$. This shows that (b_{i+1}) and (c_{i+1}) hold. Finally, since $F_i \subseteq H'_i = H_i - H_i[U_{i+1}]$ we have that $H_{i+1}[U_{i+2}] = H_i[U_{i+2}] = H'[U_{i+2}]$, and therefore (d_{i+1}) holds.

At the end of this process, we have obtained $H_t \subseteq H'[U_t]$ such that $H' - H_t$ has a $C_\ell^{(k)}$ -decomposition.

Step 3: Finish. Since H' and $H' - H_t$ are $C_\ell^{(k)}$ -divisible, we deduce $H_t \subseteq H'[U_t]$ is $C_\ell^{(k)}$ -divisible. Therefore, $H_t \in \mathcal{L}$, and by construction we know that $H_t \cup A$ has a $C_\ell^{(k)}$ -decomposition. Thus H is the edge-disjoint union of $H_t \cup A$ and $H' - H_t$ and both of them have $C_\ell^{(k)}$ -decompositions, so we deduce H has a $C_\ell^{(k)}$ -decomposition as well. \blacksquare

5. VORTEX LEMMA

We prove Lemma 4.1 by selecting subsets at random (cf. [2]).

Proof. Let $n_0 = n$ and $n_i = \lfloor \xi n_{i-1} \rfloor$ for all $i \geq 1$. In particular, note $n_i \leq \xi^i n$. Let t be the largest i such that $n_i \geq m'$ and let $m = n_{\ell+1}$. Note that $\lfloor \xi m' \rfloor \leq m \leq m'$.

Let $\xi_0 = 0$ and, for all $i \geq 1$, define $\xi_i = \xi_{i-1} + 2(\xi^i n)^{-1/3}$. Thus we have

$$\xi_{t+1} = 2n^{-1/3} \sum_{i \in [t]} (\xi^{-1/3})^i \leq 2n^{-1/3} \sum_{i \in \mathbb{N}} (\xi^{-1/3})^i \leq \frac{2(n\xi)^{-1/3}}{1 - \xi^{-1/3}} \leq \xi,$$

where in the last inequality we used $1/m' \ll \xi$ and $n \geq m'$.

Clearly, taking $U_0 = V(H)$ is a $(\delta - \xi_0, \xi, n_0)$ -vortex in H . Suppose now we have already found a $(\delta - \xi_{i-1}, \xi, n_{i-1})$ -vortex $U_0 \supseteq \cdots \supseteq U_{i-1}$ in H for some $i \leq t+1$. In particular, $\delta^{(2)}(H[U_i]) \geq (\delta - \xi_{i-1})|U_i|$. Let $U_i \subseteq U_{i-1}$ be a random subset of size n_i . By standard concentration inequalities, with positive probability we have $\delta^{(2)}(H[U_i]) \geq (\delta - \xi_{i-1} - n_i^{-1/3})|U_i|$ and $\delta^{(2)}(H[U_{i-1}], U_i) \geq (\delta - \xi_{i-1} - n_i^{-1/3})|U_i|$. Since $\xi_{i-1} + n_i^{-1/3} \leq \xi_i$, we have found a $(\delta - \xi_i, \xi, n_i)$ -vortex for H . In the end, we have found a $(\delta - \xi_{t+1}, \xi, n_{t+1})$ -vortex for H . Since $m = n_{t+1}$ and $\xi_{t+1} \leq \xi$, we are done. \blacksquare

6. TRANSFORMERS I: GADGETS

In this and the next two sections we prove Lemma 4.3, the Absorber lemma. Following Lemma 3.3, it is enough to find transformers instead of absorbers. In this part, we introduce *gadgets*, which will be building blocks of our transformers.

A k -uniform trail is a sequence of (possibly repeated) vertices such that any k consecutive vertices form an edge, and no edge appears more than once. A k -uniform tour is a k -uniform trail $v_1 \cdots v_t$ such that $v_i = v_{t-k+1+i}$ for $i \in [k-1]$. Let H be a k -graph. A *tour-trail decomposition* \mathcal{T} of H is an edge-decomposition of H into tours and trails. Note that every k -graph has a tour-trail decomposition, namely, considering each edge of H as a trail (by giving to it an arbitrary ordering). A *tour decomposition* is a tour-trail decomposition consisting only of tours. When it comes to the construction of absorbers, it is of great help to work with remainder subgraphs which admit tour decompositions. Indeed, it is straightforward to find edge-bijective homomorphisms between tours and cycles.

To construct absorbers, we will prove that actually any $C_\ell^{(k)}$ -divisible k -graph can be augmented to a new, not-so-large, subgraph which does have such a tour decomposition. This will be done in Section 7, see Lemma 7.1. In this section, we will describe certain small subgraphs which we will call *gadgets*. The augmented subgraph which we mentioned will be built as an edge-disjoint union of gadgets.

6.1. Residual graphs. Consider $k \in \mathbb{N}$ to be fixed. Now we introduce the terminology we need to describe the gadgets. Let $P = v_1 v_2 \cdots v_t$ be a trail. We define the *ends of P* to be the ordered $(k-1)$ -tuples $v_{k-1} v_{k-2} \cdots v_1$ and $v_{t-k+2} v_{t-k+3} \cdots v_t$. We denote $D(P)$ be the set of ends of P . Let \mathcal{T} be a tour-trail decomposition on vertex set V . We define the *residual di - $(k-1)$ -graph $D(\mathcal{T})$* of \mathcal{T} to be the multiset $\{D(P) : P \text{ is a trail in } \mathcal{T}\}$. Thus $D(\mathcal{T})$ consists of ordered $(k-1)$ -tuples of vertices in V , possibly counted with repetitions.

For $i \in [k-1]$ and a vertex $v \in V$, let $p_{\mathcal{T},i}(v)$ be the number of ordered $(k-1)$ -tuples in $D(\mathcal{T})$ with v being the i th vertex. We say that \mathcal{T} is *balanced* if, for all $v \in V$ and $i \in [k-1]$,

$$(6.1) \quad p_{\mathcal{T},i}(v) = p_{\mathcal{T},k-i}(v).$$

We omit \mathcal{T} from the subscript if it is known from the context.

Observe that if $v_1 \cdots v_{k-1}, v_{k-1} \cdots v_1 \in D(\mathcal{T})$, then there are trails $P_i, P_j \in \mathcal{T}$ that can be merged into a trail (if $i \neq j$) or tour (if $i = j$) with edge set $E(P_i \cup P_j)$. Thus there is another tour-trail decomposition with less trails than \mathcal{T} , which is obtained from \mathcal{T} by removing P_i, P_j and adding the tour or trail born from joining P_i and P_j . We will abuse the notation by calling the resulting tour-trail decomposition by \mathcal{T} . This merging procedure will be indicated by

$$D(\mathcal{T}) = D(\mathcal{T}) \setminus \{v_1 \cdots v_{k-1}, v_{k-1} \cdots v_1\}.$$

Given a vertex $x \in V$ and a k -tuple $\mathbf{y} = y_1 \cdots y_k$, for every $i \in [k]$ we define

$$r_i(\mathbf{y}, x) = y_1 \cdots y_{i-1} x y_{i+1} \cdots y_k, \text{ and } s_i(\mathbf{y}) = \{y_1 \cdots y_{i-1} y_{i+1} \cdots y_k\}.$$

In words, $r_i(\mathbf{y}, x)$ replaces the i th vertex in \mathbf{y} with the vertex x , whilst s_i simply skips the i th vertex of \mathbf{y} . Moreover, we define the *reverse* of \mathbf{y} as $\mathbf{y}^{-1} = y_k \cdots y_1$. Finally, given a permutation σ of $[k]$ we write $\sigma(\mathbf{y})$ for the tuple $y_{\sigma(1)} \cdots y_{\sigma(k)}$.

Given vertices $x, x' \in V$ and $(k-1)$ -tuples $\mathbf{y} = y_1 \cdots y_{k-1}$ and $\mathbf{y}' = y'_1 \cdots y'_{k-1}$, we define the following sets of $(k-1)$ -tuples:

$$\begin{aligned} \bar{S}_i^{(k-1)}(\mathbf{y}, x, x') &= \{y_1 \cdots y_{i-1} x y_{i+1} \cdots y_{k-1}, y_{k-1} \cdots y_{i+1} x' y_{i-1} \cdots y_1\} \\ &= \{r_i(\mathbf{y}, x), (r_i(\mathbf{y}, x'))^{-1}\} \text{ and} \\ \bar{T}_i^{(k-1)}(\mathbf{y}, \mathbf{y}', x, x') &= \bar{S}_j^{(k-1)}(\mathbf{y}, x, x') \cup \bar{S}_j^{(k-1)}(\mathbf{y}', x', x) \\ &= \{r_i(\mathbf{y}, x), r_i(\mathbf{y}', x'), (r_i(\mathbf{y}, x'))^{-1}, (r_i(\mathbf{y}', x'))^{-1}\}. \end{aligned}$$

For a k -tuple \mathbf{y} (instead of a $(k-1)$ -tuple), it will be convenient to consider the set $\bar{S}_i^{(k-1)}(\mathbf{y}, x, x')$ with the same definition but omitting the last element of the tuple y_k . More precisely, given a k -tuple \mathbf{y} we define

$$\bar{S}_i^{(k-1)}(\mathbf{y}, x, x') = \bar{S}_i^{(k-1)}(s_k(\mathbf{y}), x, x') = \{r_i(s_k(\mathbf{y}), x), r_i(s_k(\mathbf{y}), x')^{-1}\}.$$

An analogous definition holds for $\bar{T}_i^{(k-1)}(\mathbf{y}, \mathbf{y}', x, x')$, where \mathbf{y} and \mathbf{y}' are k -tuples. Since k will be always clear from the context, we will omit it in the notation.

The two types of $C_\ell^{(k)}$ -decomposable k -graphs to be constructed will be $B_j(x, x')$ in Corollary 6.3 (which we call ‘*balancer gadgets*’) and $T_j(\mathbf{y}, \mathbf{y}', x, x')$ in Lemma 6.6 (which we call ‘*swapper gadgets*’). This last swapper gadget has a trail-tour decomposition whose residual digraph is exactly $\bar{T}_j(\mathbf{y}, \mathbf{y}', x, x')$. Essentially, the main properties of the gadgets are:

- The rôle of balancer gadgets $B_j(x, x')$ is to enable us to adjust $p_j(x) - p_{k-1-j}(x)$ without affecting other vertices in $V(H) \setminus \{x, x'\}$. Hence, by adding edge-disjoint copies of balancer gadgets, the resulting tour-trail decomposition \mathcal{T} will be balanced (see Lemma 7.3).
- Suppose now \mathcal{T} is balanced. Consider $x \in V(H)$ and $1 \leq j \leq k/2$, we can now pair the members of $D(\mathcal{T})$ containing x into pairs $(\mathbf{y}, \mathbf{y}')$ such that x is the j th vertex in \mathbf{y} and $(k-1-j)$ th vertex in \mathbf{y}' . This is possible, as \mathcal{T} is balanced. The swapper gadget $T_j(\mathbf{y}, \mathbf{y}', x, x')$ will enable us to ‘replace’ x with a new vertex x' in both \mathbf{y} and \mathbf{y}' . By repeated applications of this gadget, this will allow us to convert \mathcal{T} into a tour decomposition (see Lemmas 7.6 and 7.10).

Throughout this section, given a tight cycle $C = v_1 \cdots v_\ell$, we often consider a trail decomposition of C consisting of two trails $v_{\sigma(1)} \cdots v_{\sigma(k)}$ and $C - v_1 \cdots v_k = v_2 \cdots v_\ell v_1 \cdots v_{k-1}$ for some permutation σ of $[k]$. Given a trail decomposition $\mathcal{T} = \{P_1, \dots, P_s\}$, for ease of checking, we often write $D(\mathcal{T}) = D(P_1) \cup \cdots \cup D(P_s)$ before merging some of the trails in \mathcal{T} , that is, deleting pairs in $D(\mathcal{T})$ of form $\{\mathbf{y}, \mathbf{y}^{-1}\}$.

The next lemma finds many trails of prescribed length which connect any given pair of ordered $(k-1)$ -tuples. It follows from [11, Lemma 2.3].

Lemma 6.1. *Let $1/n \ll \rho \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k$. Let H be a k -graph on n vertices with $\delta^{(2)}(H) \geq \varepsilon n$. Then, for every two $(k-1)$ -tuples $\mathbf{x} = v_1 \cdots v_{k-1}$ and $\mathbf{y} = v_{\ell+1} \cdots v_{\ell+k-1}$, H contains at least $\rho n^{\ell-k+1}$ trails $v_1 \cdots v_{\ell+k-1}$ on ℓ edges, each of them with no repeated vertices, except possibly those already repeated in \mathbf{x} and \mathbf{y} . In particular, if \mathbf{x}, \mathbf{y} are disjoint, then each of these trails is a tight path of length ℓ .*

6.2. Basic gadgets. We now construct a $C_\ell^{(k)}$ -decomposable k -graph, which will be a basic building block of all of the next gadgets.

Lemma 6.2. *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(2)}(H) \geq \varepsilon n$. Let $j \in [k-1]$. Let $x, x' \in V(H)$ be distinct vertices and \mathbf{y} be k -tuple of $V(H)$ such that $Y = \{y_i : i \in [k] \setminus \{j\}\} \in N_H(x) \cap N_H(x')$. Then*

there exists a $C_\ell^{(k)}$ -decomposable k -graph $G = G_j(\mathbf{y}, x, x')$ in H with $|G| = 2\ell$ and a tour-trail decomposition \mathcal{T}_j of G satisfying

$$D(\mathcal{T}_j) = \vec{S}_j(\mathbf{y}, x, x') \cup \vec{S}_1(\sigma_1(\mathbf{y}), x', x) \cup \vec{S}_{j-1}(\sigma_2(\mathbf{y}), x, x'),$$

where $\sigma_1 = j12 \cdots (j-1)(j+1) \cdots k$ and $\sigma_2 = 2 \cdots k1$. Moreover, $G[\{x, x'\} \cup Y] = \{x \cup Y, x' \cup Y\}$.

Proof. Orient $Y \cup x$ and $Y \cup x'$ into $y_k \cdots y_{j+1}xy_{j-1} \cdots y_1$ and $x'y_1 \cdots y_{j-1}y_{j+1} \cdots y_k$. By Lemma 6.1, there exist two tight cycles of length ℓ ,

$$C_1 = y_k \cdots y_{j+1}xy_{j-1} \cdots y_1 v_{k+1} \cdots v_\ell,$$

$$C_2 = x'y_1 \cdots y_{j-1}y_{j+1} \cdots y_k u_{k+1} \cdots v_\ell,$$

where v_i and u_i for $k+1 \leq i \leq \ell$ are new distinct vertices. Let $G = C_1 \cup C_2$. Define the tour-trail decomposition \mathcal{T} of $C_1 \cup C_2$ to be

$$\mathcal{T}_j = \left\{ \begin{array}{l} xy_1 \cdots y_{j-1}y_{j+1} \cdots y_k, \\ y_{k-1} \cdots y_{j+1}xy_{j-1} \cdots y_1 v_{k+1} \cdots v_\ell y_k \cdots y_{j+1}xy_{j-1} \cdots y_2, \\ y_k \cdots y_{j+1}x'y_{j-1} \cdots y_1, \\ y_1 \cdots y_{j-1}y_{j+1} \cdots y_k u_{k+1} \cdots u_\ell x'y_1 \cdots y_{j-1}y_{j+1} \cdots y_{k-1} \end{array} \right\}.$$

Hence, $D(\mathcal{T}_j)$ consists of

$$\begin{aligned} D(\mathcal{T}_j) &= \left\{ \begin{array}{ll} y_{k-1} \cdots y_{j+1}y_{j-1} \cdots y_1 x, & y_1 \cdots y_{j-1}y_{j+1} \cdots y_k, \\ y_1 \cdots y_{j-1}xy_{j+1} \cdots y_{k-1}, & y_k \cdots y_{j+1}xy_{j-1} \cdots y_2, \\ y_2 \cdots y_{j-1}x'y_{j+1} \cdots y_k, & y_{k-1} \cdots y_{j+1}x'y_{j-1} \cdots y_1, \\ y_k \cdots y_{j+1}y_{j-1} \cdots y_1, & x'y_1 \cdots y_{j-1}y_{j+1} \cdots y_{k-1} \end{array} \right\} \\ &= \vec{S}_j(\mathbf{y}, x, x') \cup \vec{S}_1(\sigma_1(\mathbf{y}), x', x) \cup \vec{S}_j(\sigma_2(\mathbf{y}), x, x'), \end{aligned}$$

where $\sigma_1 = j1 \cdots (j-1)(j+1) \cdots k$ and $\sigma_2 = 2 \cdots k1$. \blacksquare

6.3. Balancer gadgets. Next, we will use Lemma 6.2 to construct a balancer gadget $B_j = B_j(x, x')$. As mentioned before, the main property of $B_j(x, x')$ is to enable us to increase $p_{\mathcal{T},i}(x) - p_{\mathcal{T},k-1-i}(x)$ (and decreasing $p_{\mathcal{T},1}(x) - p_{\mathcal{T},k-1}(x)$) without affecting other vertices in $V \setminus \{x, x'\}$.

Corollary 6.3 (Balancer gadgets). *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(2)}(H) \geq \varepsilon n$. Let $j \in [k-1] \setminus \{1\}$ and let $x, x' \in V(H)$ be distinct. Then there exists a $C_\ell^{(k)}$ -decomposable k -graph $B_j = B_j(x, x')$ with $|B_j| = 2(j-1)\ell$ and a tour-trail decomposition \mathcal{T}'_j of B_j such that for all $i \in [k-1]$, $p_{\mathcal{T}'_j,i}(v) - p_{\mathcal{T}'_j,k-i}(v) = 0$ for all $v \in V(H) \setminus \{x, x'\}$ and*

$$p_{\mathcal{T}'_j,i}(x) - p_{\mathcal{T}'_j,k-i}(x) = p_{\mathcal{T}'_j,k-i}(x') - p_{\mathcal{T}'_j,i}(x') = \mathbb{1}_{i=j} - \mathbb{1}_{i=k-j} - j(\mathbb{1}_{i=1} - \mathbb{1}_{i=k-1}).$$

Moreover, when k is even and $j = k/2$, $p_{\mathcal{T}'_j,k/2}(v) \equiv \mathbb{1}_{v \in \{x, x'\}} \pmod{2}$.

Proof. We will proceed by induction on j . Let \mathbf{y} be a k -tuple such that $Y = \{y_i : i \in [k] \setminus \{j\}\} \in N(x) \cap N(x')$. By Lemma 6.2, there is a $C_\ell^{(k)}$ -decomposable k -graph $G_j = G_j(\mathbf{y}, x, x')$ such that

- (i) $G_j[\{x, x'\} \cup Y] = \{x \cup Y, x' \cup Y\}$,
- (ii) $|G_j| = 2\ell$, and
- (iii) there exists a tour-trail decomposition \mathcal{T}_j of G_j such that

$$D(\mathcal{T}_j) = \vec{S}_j(\mathbf{y}, x, x') \cup \vec{S}_1(\sigma_1(\mathbf{y}), x', x) \cup \vec{S}_{j-1}(\sigma_2(\mathbf{y}), x, x'),$$

where $\sigma_1 = j12 \cdots (j-1)(j+1) \cdots k$ and $\sigma_2 = 2 \cdots k1$.

Note that, for all $i \in [k-1]$ and $v \in V(H) \setminus \{x, x'\}$, we have $p_{\mathcal{T}_j,i}(v) - p_{\mathcal{T}_j,k-i}(v) = 0$ as each of $\vec{S}_j(\mathbf{y}, x, x')$, $\vec{S}_1(\sigma_1(\mathbf{y}), x', x)$ and $\vec{S}_{j-1}(\sigma_2(\mathbf{y}), x, x')$ contributes zero. Moreover,

$$p_{\mathcal{T}_j,i}(x) - p_{\mathcal{T}_j,k-i}(x) = p_{\mathcal{T}_j,k-i}(x') - p_{\mathcal{T}_j,i}(x')$$

$$(6.2) \quad = (\mathbb{1}_{i=j} - \mathbb{1}_{i=k-j}) - (\mathbb{1}_{i=1} - \mathbb{1}_{i=k-1}) - (\mathbb{1}_{i=j-1} - \mathbb{1}_{i=k-j+1}).$$

For $j = 2$, we set $B_2 = G_2$ and we are done. For $j > 2$, there exists $B_{j-1}(x, x')$ edge-disjoint from G_j , by our induction hypothesis. Let \mathcal{T}'_{j-1} be the corresponding tour-trail decomposition. Set $B_j = G_j \cup B_{j-1}(x, x')$. Clearly $|B_j| = |G_j| + |B_{j-1}| = 2(j-1)\ell$. Note that B_j is $C_\ell^{(k)}$ -decomposable and has a tour-trail decomposition $\mathcal{T}'_j = \mathcal{T}_j \cup \mathcal{T}'_{j-1}$. Together with (6.2), we deduce that \mathcal{T}'_j satisfies the desired properties. The moreover statement can be verified similarly. \blacksquare

6.4. Swapper gadgets. The construction of swapper gadgets requires more steps. We start with the following proposition.

Proposition 6.4. *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq \varepsilon n$. Let $x, x' \in V(H)$ be distinct vertices and \mathbf{y} be a $(k-1)$ -tuple of $V(H)$ such that $\{x, x'\} \cup \{y_i : 2 \leq i \leq k-1\}$ is of size k . Then, there exists a vertex $y_k \in V(H)$, a $C_\ell^{(k)}$ -decomposable k -graph $F_1 = F_1(\mathbf{y}, x, x')$ in H with $|F_1| = 3\ell$ and a tour-trail decomposition \mathcal{T}_1 such that*

$$D(\mathcal{T}_1) = \begin{cases} \bar{S}_1(\mathbf{y}, x, x') \cup \{xx', xx'\} & \text{if } k = 3, \\ \bar{S}_1(\mathbf{y}, x, x') \cup \{xx'y_k \dots y_4, y_4 \dots y_k xx'\} & \text{if } k \geq 4. \end{cases}$$

Moreover, $F_1[\{x, x', y_2, \dots, y_{k-1}\}] = \emptyset$.

Proof. Let $y_k \in N(xy_2 \dots y_{k-1}) \cap N(x'y_2 \dots y_{k-1}) \cap N(xx'y_3 \dots y_{k-1})$, which exists by our assumption (here for $k = 3$ we consider $y_3 \dots y_2$ to be empty). By Lemma 6.1, there exist three tight cycles of length ℓ

$$\begin{aligned} C_1 &= y_2 \dots y_k x' u_{k+1} \dots u_\ell, \\ C_2 &= xy_2 \dots y_k v_{k+1} \dots v_\ell, \text{ and} \\ C_3 &= y_3 \dots y_k x' w_{k+1} \dots w_\ell, \end{aligned}$$

where u_i, v_i, w_i are all distinct new vertices. Let $T_1 = C_1 \cup C_2 \cup C_3$. Consider the trail decomposition \mathcal{T}_1 of T_1 such that

$$\mathcal{T}_1 = \left\{ \begin{array}{l} y_3 \dots y_k x' u_{k+1} \dots u_\ell y_2 \dots y_k, \\ x' y_2 \dots y_k, \\ y_2 \dots y_k v_{k+1} \dots v_\ell x y_2 \dots y_{k-1}, \\ y_2 \dots y_k x, \\ y_4 \dots y_k x' w_{k+1} \dots w_\ell y_3 \dots y_k x', \\ y_3 \dots y_k x x' \end{array} \right\}$$

and so

$$D(\mathcal{T}_1) = \left\{ \begin{array}{ll} \overline{x' y_k \dots y_3}, & \overline{y_2 \dots y_k}, \\ \overline{y_{k-1} \dots y_2 x'}, & \overline{y_2 \dots y_k}, \\ \overline{y_k \dots y_2} & \overline{xy_2 \dots y_{k-1}}, \\ \overline{y_k \dots y_2}, & \overline{y_3 \dots y_k x}, \\ \overline{xx' y_k \dots y_4}, & \overline{y_3 \dots y_k x'}, \\ \overline{xy_k \dots y_3}, & \overline{y_4 \dots y_k x x'} \end{array} \right\}$$

as required. \blacksquare

We construct a swapper gadget of the form $T_1(\mathbf{y}, \mathbf{y}', x, x')$ in the next proposition.

Proposition 6.5 (Swapper gadget – case $j = 1$). *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ such that $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq \varepsilon n$. Consider vertices $x, x' \in V(H)$ and $(k-1)$ -tuples \mathbf{y}, \mathbf{y}' of $V(H)$ such that both $\{x, x'\} \cup \{y_i : 2 \leq i \leq k-1\}$ and $\{x, x'\} \cup \{y'_i : 2 \leq i \leq k-1\}$ are of size k . Then there exists a $C_\ell^{(k)}$ -decomposable k -graph $T_1 = T_1(\mathbf{y}, \mathbf{y}', x, x')$ in H a tour-trail decomposition \mathcal{T}_1 of T_1 such that*

- (i) $T_1[\{x, x'\} \cup \{y_1, \dots, y_{k-1}\} \cup \{y'_1, \dots, y'_{k-1}\}] = \emptyset$ and $|T_1| = 2\ell k$ and

$$(ii) D(\mathcal{T}_1) = \bar{T}_1(\mathbf{y}, \mathbf{y}', x, x').$$

Proof. We apply Proposition 6.4 twice, the first time for x, x' and \mathbf{y} as input and the second time with x', x and \mathbf{y}' as input (we exchange the rôles of x and x'). This yields vertices $y_k, y'_k \in V(H)$ and two $C_\ell^{(k)}$ -decomposable k -graphs $F = F_1(\mathbf{y}, x, x')$ and $F' = F_1(\mathbf{y}', x', x)$ such that

- (a₁) $V(F) \cap V(F') \subseteq \{x, x'\} \cup \{y_i, y'_i : i \in [k-1]\}$ and $|F| = |F'| = 3\ell$ and
- (a₂) there exists a tour-trail decomposition \mathcal{T} of $F \cup F'$ such that

$$D(\mathcal{T}) = T_1(\mathbf{y}, \mathbf{y}', x', x) \cup \{xx'y_k \cdots y_4, y_4 \cdots y_k x x', x'x y'_k \cdots y'_4, y'_4 \cdots y'_k x'x\},$$

where, for $k = 3$, we interpret the strings $y_k \cdots y_4$ and $y'_k \cdots y'_4$ to be empty.

If $k = 3$, then $D(\mathcal{T}) = T_1(\mathbf{y}, \mathbf{y}', x', x) \cup \{x'x, x'x, xx', xx'\} = T_1(\mathbf{y}, \mathbf{y}', x', x)$, thus we are done. So we may assume that $k \geq 4$. Note that if we have $y_i = y'_i$ for all $i \in \{4, \dots, k\}$, then $D(\mathcal{T})$ is as desired. Thus, our aim is to ‘replace’ y_i, y'_i with a new vertex z_i , for each $i \in \{4, \dots, k\}$. We do this in turns, as follows. For each $i \in \{4, \dots, k\}$, let

$$z_i \in N(xx'z_4 \cdots z_{i-1}y_i \cdots y_k) \cap N(xx'z_4 \cdots z_{i-1}y'_i \cdots y'_k)$$

be a new vertex (here we consider $z_4 \cdots z_3$ to be empty). Consider the two ordered edges $z_i \cdots z_4 xx' y_k \cdots y_i$ and $z_i \cdots z_4 x' y'_k \cdots y'_i$ and apply Lemma 6.1 to obtain two tight cycles of length ℓ

$$C^i = z_i \cdots z_4 xx' y_k \cdots y_i v_{k+1}^i \cdots v_\ell^i \text{ and } D^i = z_i \cdots z_4 x' y'_k \cdots y'_i w_{k+1}^i \cdots w_\ell^i$$

such that v_j^i, w_j^i are new vertices. Define a tour-trail decomposition \mathcal{T}^i of $C^i \cup D^i$ such that

$$\mathcal{T}^i = \left\{ \begin{array}{l} z_{i-1} \cdots z_4 xx' y_k \cdots y_i v_{k+1}^i \cdots v_\ell^i z_i \cdots z_4 xx' y_k \cdots y_{i+1}, \\ z_i \cdots z_4 x' y'_k \cdots y'_i, \\ z_{i-1} \cdots z_4 x' y'_k \cdots y'_i w_{k+1}^i \cdots w_\ell^i z_i \cdots z_4 x' y'_k \cdots y'_{i+1}, \\ z_i \cdots z_4 xx' y'_k \cdots y'_i \end{array} \right\}.$$

Note that

$$\begin{aligned} D(\mathcal{T}^i) &= \left\{ \begin{array}{ll} y_i \cdots y_k x' x z_4 \cdots z_{i-1}, & z_i \cdots z_4 xx' y_k \cdots y_{i+1}, \\ y_{i+1} \cdots y_k xx' z_4 \cdots z_i, & z_{i-1} \cdots z_4 x' y'_k \cdots y'_i, \\ y'_i \cdots y'_k xx' z_4 \cdots z_{i-1}, & z_i \cdots z_4 x' y'_k \cdots y'_{i+1}, \\ y'_{i+1} \cdots y'_k x' z_4 \cdots z_i, & z_{i-1} \cdots z_4 xx' y'_k \cdots y'_i \end{array} \right\} \\ &= \left\{ \begin{array}{l} y_i \cdots y_k x' x z_4 \cdots z_{i-1}, \\ y'_i \cdots y'_k xx' z_4 \cdots z_{i-1}, \\ z_{i-1} \cdots z_4 x' y'_k \cdots y'_i, \\ z_{i-1} \cdots z_4 xx' y'_k \cdots y'_i \end{array} \right\} \cup \left\{ \begin{array}{l} z_i \cdots z_4 xx' y_k \cdots y_{i+1}, \\ z_i \cdots z_4 x' y'_k \cdots y'_{i+1}, \\ y_{i+1} \cdots y_k xx' z_4 \cdots z_i, \\ y'_{i+1} \cdots y'_k x' z_4 \cdots z_i \end{array} \right\}. \end{aligned}$$

When $i = k$, then the second set can be simplified to an empty set. Note that $D(\mathcal{T}^4), \dots, D(\mathcal{T}^k)$ forms a ‘telescoping’ set of residual di- $(k-1)$ -graphs, so we deduce that

$$D\left(\bigcup_{4 \leq i \leq k} \mathcal{T}^i\right) = \{xx'y'_k \cdots y'_4, y'_4 \cdots y'_k xx', x'xy_k \cdots y_4, y_4 \cdots y_k x'x\}.$$

We are done by setting $T_1 = F \cup F' \cup \bigcup_{4 \leq i \leq k} (C^i \cup D^i)$ and $\mathcal{T}_1 = \mathcal{T} \cup \bigcup_{4 \leq i \leq k} \mathcal{T}^i$. ■

We can now describe the general version of the swapper gadget T_j , for all $j \in [k-1]$.

Lemma 6.6 (Swapper gadget – general case). *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ such that $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq \varepsilon n$. Let $j \in [k-1]$ and consider distinct vertices $x, x' \in V(H)$ and $(k-1)$ -tuples \mathbf{y}, \mathbf{y}' of $V(H)$ such that both $\{x, x'\} \cup \{y_i : i \in [k-1] \setminus \{j\}\}$ and $\{x, x'\} \cup \{y'_i : i \in [k-1] \setminus \{j\}\}$ are of size k . Then there exists a $C_\ell^{(k)}$ -decomposable k -graph $T_j = T_j(\mathbf{y}, \mathbf{y}', x, x')$ and a tour-trail decomposition \mathcal{T}_j of T_j such that*

- (i) $T_j[\{x, x'\} \cup \{y_1, \dots, y_{k-1}\} \cup \{y'_1, \dots, y'_{k-1}\}] = \emptyset$ and $|T_j| \leq 3^j \ell k$ and

$$(ii) \quad D(\mathcal{T}_j) = \bar{T}_j(\mathbf{y}, \mathbf{y}', x, x').$$

Proof. We proceed by induction on j . Note that Proposition 6.5 implies the case when $j = 1$, so we may assume that $j \geq 2$. Let

$$y_k \in N(xy_1 \dots y_{j-1} y_{j+1} \dots y_{k-1}) \cap N(x'y_1 \dots y_{j-1} y_{j+1} \dots y_{k-1})$$

be a new vertex. By Lemma 6.2, there exists a $C_\ell^{(k)}$ -decomposable k -graph $G_j = G_j(\mathbf{y}, x, x')$ with $|G_j| = 2\ell$ and a tour-trail decomposition \mathcal{G}_j of G_j satisfying

$$D(\mathcal{G}_j) = \bar{S}_j(\mathbf{y}, x, x') \cup \bar{S}_1(\sigma_1(\mathbf{y}), x', x) \cup \bar{S}_{j-1}(\sigma_2(\mathbf{y}), x', x),$$

where $\sigma_1 = j12 \dots (j-1)(j+1) \dots k$ and $\sigma_2 = 2 \dots k1$. Analogously, there is a $C_\ell^{(k)}$ -decomposable k -graph $G'_j = G'_j(\mathbf{y}', x', x)$ with $|G'_j| = 2\ell$ and a tour-trail decomposition \mathcal{G}'_j of G'_j satisfying

$$D(\mathcal{G}'_j) = \bar{S}_j(\mathbf{y}', x', x) \cup \bar{S}_1(\sigma_1(\mathbf{y}'), x, x') \cup \bar{S}_{j-1}(\sigma_2(\mathbf{y}'), x, x').$$

Note that

$$(6.3) \quad |G_j \cup G'_j| = 4\ell$$

and

$$(6.4) \quad D(\mathcal{G}_j) \cup D(\mathcal{G}'_j) = \bar{T}_j(\mathbf{y}, \mathbf{y}', x, x') \cup \bar{T}_1(\sigma_1(\mathbf{y}), \sigma_1(\mathbf{y}'), x, x') \cup \bar{T}_{j-1}(\sigma_2(\mathbf{y}), \sigma_2(\mathbf{y}'), x, x').$$

Due to the induction hypothesis, there are $C_\ell^{(k)}$ -decomposable k -graphs

$$T_1 = T_1(\sigma_1(\mathbf{y}), \sigma_1(\mathbf{y}'), x', x) \quad \text{and} \quad T_{j-1} = T_{j-1}(\sigma_2(\mathbf{y}), \sigma_2(\mathbf{y}'), x', x)$$

and a tour-trail decomposition \mathcal{T}_j^* of $T_1 \cup T_{j-1}$ such that their union $T_j^* = T_1 \cup T_{j-1}$ satisfies

$$(i') \quad T_j^*[\{x, x'\} \cup \{y_1, \dots, y_{k-1}\} \cup \{y_1, \dots, y_{k-1}\}] = \emptyset \quad \text{and} \quad |T_j^*| \leq 2 \cdot 3^{j-1} \ell k,$$

$$(ii') \quad D(\mathcal{T}_j^*) = \bar{T}_1(\sigma_1(\mathbf{y}), \sigma_1(\mathbf{y}'), x', x) \cup \bar{T}_{j-1}(\sigma_2(\mathbf{y}), \sigma_2(\mathbf{y}'), x', x)$$

We set $T_j = G_j \cup G'_j \cup T_j^*$ and $\mathcal{T}_j = \mathcal{G}_j \cup \mathcal{G}'_j \cup \mathcal{T}_j^*$. By (6.3) and (i'), we deduce that

$$|T_j| \leq 4\ell + 2 \cdot 3^{j-1} \ell k \leq 3^j \ell k.$$

Moreover (6.4) and (ii') imply that $D(\mathcal{T}_j) = \bar{T}_j(\mathbf{y}, \mathbf{y}', x, x')$ as required. \blacksquare

7. TRANSFORMERS II: TOUR-TRAIL DECOMPOSITIONS

Here, we use the gadgets constructed in the previous section to prove that any $C_\ell^{(k)}$ -divisible k -graph can be augmented to a new, not-too-large, subgraph which has a tour decomposition. That is the content of the next crucial lemma, whose proof will be given at the end of this section. Note that we only require $\deg_G(v)$ is divisible by k for all vertices $v \in V(G)$ instead of $C_\ell^{(k)}$ -divisible.

Lemma 7.1. *Let $1/n \ll \varepsilon \ll \rho, 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq \varepsilon n$. Let G be a k -graph with $V(G) \subseteq V(H)$ and $m = |V(G)| \leq \varepsilon n^{1/k(k+1)}$ such that $\deg_G(v)$ is divisible by k for all $v \in V(G)$. Then $H - G$ contains a $C_\ell^{(k)}$ -decomposable subgraph J such that $G \cup J$ has a tour decomposition, $J[V(G)] = \emptyset$ and $|G \cup J| \leq 3^{k+2} k^4 \ell^2 m^{k+1}$. Moreover, if G' has an edge-bijective homomorphism to G with $V(G') \subseteq V(H) \setminus V(G)$, then we have $H - G - G'$ contains a subgraph J' such that $G' \cup J'$ is edge-bijective homomorphic to $G \cup J$, $J'[V(G')] = \emptyset$ and $V(G \cup J) \cap V(G' \cup J') = \emptyset$.*

As described before, the lemma will be proven by starting with any tour-trail decomposition of G , and adding gadgets to it repeatedly. We will first use balancer gadgets to make sure we have a balanced tour-trail decomposition, and then we will use swapper gadgets to eliminate any remaining trails one by one.

7.1. Basic properties. We begin by stating basic properties of any tour-trail decomposition. Recall that $p_{\mathcal{T},i}(v)$ is the number of (directed) edges of \mathcal{T} where v is the i th vertex.

Proposition 7.2. *Let H be a k -graph such that $\deg(v)$ is divisible by k for every vertex $v \in V(H)$. Let \mathcal{T} be a tour-trail decomposition of H . Then, for each $v \in V(H)$,*

$$\sum_{i \in [k-1]} ip_{\mathcal{T},i}(v) \equiv 0 \pmod{k}.$$

Moreover, if k is even and \mathcal{T} is balanced, then $p_{\mathcal{T},k/2}(v)$ is even for all $v \in V(H)$.

Proof. Note that only trails in \mathcal{T} contribute to $\sum_{i \in [k-1]} ip_{\mathcal{T},i}(x)$. Moreover, for any tour C , we have $\deg_C(v) \equiv 0 \pmod{k}$ for all $v \in V(H)$. Hence by deleting all tours in \mathcal{T} and their corresponding edges in H , we may assume that \mathcal{T} consists of trails only.

Fix $v \in V(H)$. Let $T = v_1 \dots v_t$ be a trail in \mathcal{T} and consider $I \subseteq [t]$ such that $v_i = v$. Let

$$\phi_T(v) = \sum_{i \in [t-k+1]} (\mathbb{1}_{v=v_i} + \mathbb{1}_{v=v_{i+1}} + \dots + \mathbb{1}_{v=v_{i+k-1}}).$$

Observe that every time the trail ‘pass through v ’ it increases $\phi_T(v)$ by k except if it is at the beginning or the end of T . More precisely, it is not hard to check that

$$\begin{aligned} \phi_T(v) &= k|I \cap [k, t-k]| + \sum_{i \in [k-1] \cup [t-k+1, t]} (\mathbb{1}_{v=v_i} + \dots + \mathbb{1}_{v=v_{i+k-1}}) \\ &\equiv \sum_{i \in [k-1]} ip_{i,T}(v) \pmod{k}. \end{aligned}$$

On the other hand, it is easy to see that $\phi_T(v) = \sum_{e \in T} \mathbb{1}_{v \in e}$. Thus, summing over all trails in \mathcal{T} , we get

$$0 \equiv \deg_H(v) = \sum_{T \in \mathcal{T}} \phi_T(v) \equiv \sum_{i \in [k-1]} ip_{i,\mathcal{T}}(v) \pmod{k}.$$

Furthermore, suppose that k is even and \mathcal{T} is balanced (see definition in (6.1)). Then

$$\sum_{i \in [k-1]} ip_{\mathcal{T},i}(v) = \sum_{i \in [k/2-1]} kp_{\mathcal{T},i}(v) + (k/2)p_{\mathcal{T},k/2}(v).$$

Since this is equivalent to $0 \pmod{k}$, we get that $p_{\mathcal{T},k/2}(v)$ is even. \blacksquare

7.2. Balancing. Recall that any k -graph admits a trail decomposition, by orienting edges arbitrarily. We begin by using balancer gadgets (as given by Corollary 6.3) repeatedly to obtain a balanced tour-trail decomposition of G .

Lemma 7.3. *Let $1/n \ll \varepsilon \ll \rho, 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices and $\delta^{(3)}(H) \geq \varepsilon n$. Let G be a k -graph with $V(G) \subseteq V(H)$, $m = |V(G)| \leq \varepsilon n^{1/k^2}$ and such that $\deg_G(v)$ is divisible by k for all $v \in V(G)$. Then $H - G$ contains a $C_\ell^{(k)}$ -decomposable J such that $G \cup J$ has a balanced tour-trail decomposition \mathcal{T}^* , $J[V(G)] = \emptyset$ and $|G \cup J| \leq \ell m^{k+1}$.*

Proof. Let \mathcal{T}_0 be an arbitrary tour-trail decomposition of G . The following claim forms the basis of our proof and allows us to adjust the values of $p_{\mathcal{T}_0,k/2}(v)$.

Claim 7.4. *Suppose k is even. Then $H - G$ contains a $C_\ell^{(k)}$ -divisible subgraph $J_{k/2}$ with $|J_{k/2}| \leq k\ell m/2$ and $J_{k/2}[V(G)] = \emptyset$ such that there exists a tour-trail decomposition $\mathcal{T}_{k/2}$ of $J_{k/2}$ satisfying, for all $v \in V(H)$,*

$$\begin{aligned} p_{\mathcal{T}_{k/2},k/2}(v) &\equiv \mathbb{1}_{p_{\mathcal{T}_0,k/2}(v) \text{ is odd}} \pmod{2}, \\ |p_{\mathcal{T}_{k/2},1}(v) - p_{\mathcal{T}_{k/2},k-1}(v)| &= (k/2) \mathbb{1}_{p_{\mathcal{T}_0,k/2}(v) \text{ is odd}}, \\ p_{\mathcal{T}_{k/2},i}(v) - p_{\mathcal{T}_{k/2},k-i}(v) &= 0 \text{ if } i \in [2, k-2]. \end{aligned}$$

Proof of claim. Note that $\sum_{v \in V(H)} p_{\mathcal{T}_0, k/2}(v) = |D(\mathcal{T}_0)|$ is twice the number of trails in $D(\mathcal{T}_0)$. Without loss of generality, let $v_1, \dots, v_{2s} \in V(H)$ be the vertices in $V(G)$ such that $p_{\mathcal{T}_0, k/2}(v_j)$ is odd. For each $j \in [s]$, we apply Corollary 6.3 and obtain edge-disjoint balancer gadgets $B_{k/2}(v_{2j-1}, v_{2j})$ in $H - G$. Let $J_{k/2}$ be the union of these balancer gadgets. Clearly, $|J_{k/2}| \leq sk\ell/2 \leq k\ell m/2$. Let $\mathcal{T}_{k/2}$ be the tour-trail decomposition of $J_{k/2}$, which is the union of the corresponding tour-trail decompositions of each $B_{k/2}(v_{2j-1}, v_{2j})$. For all $v \in V(H)$, we have $p_{\mathcal{T}_{k/2}, k/2}(v) \equiv 1 \pmod{2}$ if and only if $v \in \{v_1, \dots, v_{2s}\}$, which proves the first required property. We can deduce the other two properties using the properties of the corresponding tour-trail decomposition of each $B_{k/2}(v_{2j-1}, v_{2j})$. This proves the claim. \square

If k is even, then let $J_{k/2}$ be given by Claim 7.4, otherwise just set $J_{k/2} = \emptyset$. The next claim allows us to adjust the values of $p_{\mathcal{T}_0, i}(v)$ for $2 \leq i < k/2$.

Claim 7.5. *For each $2 \leq i < k/2$, $H - G - J_{k/2}$ contains a $C_\ell^{(k)}$ -divisible subgraph J_i such that there exists a tour-trail decomposition \mathcal{T}_i of J_i satisfying, for all $v \in V(H)$ and $i' \in [k-1]$,*

$$p_{\mathcal{T}_i, i'}(v) - p_{\mathcal{T}_i, k-i'}(v) = \begin{cases} p_{\mathcal{T}_0, k-i'}(v) - p_{\mathcal{T}_0, i'}(v) & \text{if } i' \in \{i, k-i\}, \\ -i(p_{\mathcal{T}_0, k-i'}(v) - p_{\mathcal{T}_0, i'}(v)) & \text{if } i' \in \{1, k-1\}, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, $|J_i| \leq 4i\ell \binom{m}{k}$, $J_i[V(G)] = \emptyset$ and the J_i 's are edge-disjoint.

Proof of claim. Let $2 \leq i < k/2$. Suppose that we have already constructed subgraphs J_2, \dots, J_{i-1} . We now construct J_i as follows. Let $H' = H - G - J_{k/2} - \bigcup_{i' \in [2, i-1]} J_{i'}$.

For all $i' \in [k-1]$, note that $\sum_{v \in V(H)} p_{\mathcal{T}_0, i'}(v) = |D(\mathcal{T}_0)|$ and so

$$\sum_{v \in V(H)} (p_{\mathcal{T}_0, i}(v) - p_{\mathcal{T}_0, k-i}(v)) = 0.$$

Define a multi-digraph H_i on $V(H)$ such that, for all $v \in V(H)$,

$$d_{H_i}^-(v) = \max\{p_{\mathcal{T}_0, i}(v) - p_{\mathcal{T}_0, k-i}(v), 0\} \text{ and } d_{H_i}^+(v) = \max\{p_{\mathcal{T}_0, k-i}(v) - p_{\mathcal{T}_0, i}(v), 0\}.$$

Note that H_i can be constructed greedily. Note that $|H_i| = |D(\mathcal{T}_0)|$ is twice the number of trails in \mathcal{T}_0 , so $|H_i| \leq 2 \binom{m}{k}$. For each directed edge $xy \in H_i$, we apply Corollary 6.3 and obtain edge-disjoint balancer gadgets $B_i(x, y)$ in H' . Let J_i be the union of these balancer gadgets. Clearly, $|J_i| \leq 2i\ell|H_i| \leq 4i\ell \binom{m}{k}$. Let \mathcal{T}_i be the tour-trail decomposition of J_i , which is the union of the corresponding tour-trail decompositions of each $B_i(x, y)$. It is straightforward to check that \mathcal{T}_i has the desired properties, which proves the claim. \square

For each $2 \leq i < k/2$, let J_i be given by Claim 7.5. Together with $J_{k/2}$, we have then edge-disjoint $J_2, \dots, J_{\lfloor k/2 \rfloor}$ for any k . Let $H^* = H - G - \bigcup_{2 \leq i \leq \lfloor k/2 \rfloor} J_i$ and set $\mathcal{T}' = \mathcal{T}_0 \cup \bigcup_{2 \leq i \leq \lfloor k/2 \rfloor} \mathcal{T}_i$. Recall that $\sum_{i \in [k-1]} p_{\mathcal{T}_0, i}(v)$ is the number of $(k-1)$ -tuples in $D(\mathcal{T}_0)$ containing v , so $\sum_{i \in [k-1]} p_{\mathcal{T}_0, i}(v) \leq 2 \binom{m}{k}$. For $i \in \{2, \dots, k-2\}$ and $v \in V(H)$, we have

$$(7.1) \quad p_{\mathcal{T}', i}(v) = p_{\mathcal{T}', k-i}(v),$$

$$(7.2) \quad p_{\mathcal{T}', k/2}(v) \equiv 0 \pmod{2} \text{ if } k \text{ is even, and}$$

$$(7.3) \quad |p_{\mathcal{T}', 1}(v) - p_{\mathcal{T}', k-1}(v)| \leq \sum_{i \in [k-1]} i p_{\mathcal{T}_0, i}(v) \leq 2(k-1) \binom{m}{k}.$$

Moreover, $p_{\mathcal{T}', 1}(v) = p_{\mathcal{T}', k-1}(v)$ for all $v \in V(H) \setminus V(G)$.

We now balance $p_{\mathcal{T}',1}(v)$ and $p_{\mathcal{T}',k-1}(v)$ as follows. Note that $\lceil k/2 \rceil - 1$ is the largest integer which is strictly less than $k/2$. We have

$$\begin{aligned} p_{\mathcal{T}',1}(v) - p_{\mathcal{T}',k-1}(v) &\stackrel{(7.1)}{=} p_{\mathcal{T}',1}(v) - p_{\mathcal{T}',k-1}(v) + \sum_{2 \leq i \leq \lceil k/2 \rceil - 1} i (p_{\mathcal{T}',i}(v) - p_{\mathcal{T}',k-i}(v)) \\ &\stackrel{(7.2)}{=} \sum_{1 \leq i \leq k-1} i p_{\mathcal{T}',i}(v) \stackrel{\text{Prop. 7.2}}{\equiv} 0 \pmod{k}. \end{aligned}$$

For each $v \in V(G)$, let

$$b(v) = (p_{\mathcal{T}',1}(v) - p_{\mathcal{T}',k-1}(v))/k,$$

so $b(v) \in \mathbb{Z}$. Define (greedily) a multi-digraph H_1 on $V(H)$ such that, for all $v \in V(H)$,

$$d_{H_1}^-(v) = \max\{b(v), 0\} \text{ and } d_{H_1}^+(v) = \max\{-b(v), 0\}.$$

By (7.3), $\Delta^\pm(H_1) \leq 2 \binom{m}{k}$ and $|H_1| \leq m \binom{m}{k}$. For each directed edge $xy \in H_1$, we apply Corollary 6.3 to obtain edge-disjoint balancer gadgets $B_{k-1}(x, y)$ in H^* . We call J_1 the union of these balancer gadgets. Clearly, $|J_1| \leq 2(k-1)\ell|H_1| \leq 2(k-1)\ell m \binom{m}{k}$.

Let $J = \bigcup_{1 \leq i \leq \lceil k/2 \rceil} J_i$. Note that

$$|G \cup J| \leq \binom{m}{k} + 2(k-1)\ell m \binom{m}{k} + \sum_{2 \leq i \leq \lceil k/2 \rceil} 4i\ell \binom{m}{k} \leq 2k\ell m \binom{m}{k} \leq \ell m^{k+1}.$$

Let \mathcal{T}_1 be the tour-trail decomposition of J_1 , which is the union of the corresponding tour-trail decompositions of each $B_{k-1}(x, y)$. Note that given a $B_{k-1}(x, y)$ and its tour-trail decomposition $\mathcal{T} = \mathcal{T}(x, y)$, we have, for all $v \in V(H)$ and $2 \leq i \leq k-2$,

$$p_{\mathcal{T},1}(v) - p_{\mathcal{T},k-1}(v) = -k(\mathbb{1}_{v=x} - \mathbb{1}_{v=y}) \text{ and } p_{\mathcal{T},i}(v) - p_{\mathcal{T},k-i}(v) = 0.$$

Hence $\mathcal{T}^* = \mathcal{T}_1 \cup \mathcal{T}'$ is a balanced tour-trail decomposition of $J \cup G$, as required. \blacksquare

7.3. Focusing. The following lemma shows that all the residual $D(\mathcal{T}^*)$ can be moved onto a set of fixed $k-1$ vertices.

Lemma 7.6. *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices and $\delta^{(3)}(H) \geq \varepsilon n$. Let G be a k -graph with $V(G) \subseteq V(H)$ and $m = |V(G)| \leq \varepsilon n^{1/k^2}/2$ such that $\deg_G(v)$ is divisible by k for all vertices $v \in V(G)$. Suppose that m is prime and $|G| < m/k$. Suppose that G has a balanced tour-trail decomposition \mathcal{T} . Let $z_1, \dots, z_{k-1} \in V(H) \setminus V(G)$ be distinct vertices. Then $H - G$ contains a $C_\ell^{(k)}$ -decomposable J^* such that $|J^*| \leq 3^k k \ell m$, $J^*[V(G)] = \emptyset$ and $G \cup J^*$ has a balanced tour-trail decomposition \mathcal{T}^* satisfying, for all $i \in [k-1]$, and $v \in V(H)$, $p_{\mathcal{T}^*,i}(v) = 0$ unless $v \in \{z_i, z_{k-i}\}$ and $|D(\mathcal{T}^*)| \leq 3m$.*

We first outline the proof of Lemma 7.6. Take $z_1, \dots, z_{k-1} \in V$ distinct vertices. Recall the definition of r_j at the beginning of Section 6. For $1 \leq i \leq \lceil k/2 \rceil$, define the functions $\zeta_i, \bar{\zeta}_i : V^{k-1} \rightarrow V^{k-1}$ be such that, for $\mathbf{a} = a_1 \cdots a_{k-1} \in V^{k-1}$,

$$\begin{aligned} \zeta_i(\mathbf{a}) &= r_{k-i}(r_i(\mathbf{a}, z_i), z_{k-i}) = a_1 \cdots a_{i-1} z_i a_{i+1} \cdots a_{k-i-1} z_{k-i} a_{k-i+1} \cdots a_{k-1} \text{ and} \\ \bar{\zeta}_i(\mathbf{a}) &= r_{k-i}(r_i(\mathbf{a}, z_{k-i}), z_i) = a_1 \cdots a_{i-1} z_{k-i} a_{i+1} \cdots a_{k-i-1} z_i a_{k-i+1} \cdots a_{k-1}. \end{aligned}$$

That is, ζ_i replaces the i th and $(k-i)$ th vertices with the vertices z_i and z_{k-i} , respectively, whereas $\bar{\zeta}_i$ replaces the i th and $(k-i)$ th vertices with vertices z_{k-i} and z_i , respectively. If k is even and $i = k/2$, then both $\zeta_{k/2}$ and $\bar{\zeta}_{k/2}$ replace the $(k/2)$ th vertex with $z_{k/2}$, this is $\zeta(\mathbf{a}) = \bar{\zeta}(\mathbf{a}) = r_{k/2}(\mathbf{a}, z_{k/2})$.

We say that a tour-trail decomposition \mathcal{T}' is an i -convert of \mathcal{T} if $D(\mathcal{T})$ can be partitioned into D_1 and D_2 of equal size such that $D(\mathcal{T}') = \zeta_i(D_1) \cup \bar{\zeta}_i(D_2)$. Note that the definitions of $\zeta_i, \bar{\zeta}_i$ and i -convert depend on a choice of z_1, \dots, z_{k-1} . However, such choice will be always clear from the context, so we omit it on the notation.

Let \mathcal{T}_0 be a balanced tour-trail decomposition of G . Our aim is to construct tour-trail decompositions $\mathcal{T}_1, \dots, \mathcal{T}_{\lfloor k/2 \rfloor}$ such that \mathcal{T}_i is an i -convert of \mathcal{T}_{i-1} . Notice that $\mathcal{T}_{\lfloor k/2 \rfloor}$ will be the desired tour-trail decomposition.

We will also need the following notation. Let \mathcal{T} be a tour-trail decomposition and $1 \leq i < k/2$. Define $A_i(\mathcal{T})$ to be the multidigraph on $V(H)$ such that every ordered tuple $v_1 \cdots v_{k-i}$ in $D(\mathcal{T})$ corresponds to a distinct directed edge $v_i v_{k-i}$ in $A_i(\mathcal{T})$.

The following is immediate from our definition of i -convert and $A_j(\mathcal{T})$.

Proposition 7.7. *Let $k \geq 3$ and $1 \leq i < k/2$. Let V be a set of vertices, and let $z_1, \dots, z_{k-1} \in V$ be distinct vertices. Let \mathcal{T} and \mathcal{T}' be tour-trail decompositions of two (not necessarily of the same) subgraphs in V . Suppose \mathcal{T}' is an i -convert of \mathcal{T} . Then $A_j(\mathcal{T}') = A_j(\mathcal{T})$ for all $1 \leq j < k/2$ such that $j \neq i$.*

The next lemma shows that we can always get a tour-trail decomposition such that $A_i(\mathcal{T})$ is strongly connected for all $1 \leq i < k/2$ and spans $V(G)$. The proof is simple and follows by greedily adding new arcs to $A_i(\mathcal{T})$.

Lemma 7.8. *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq \varepsilon n$. Let $U \subseteq V(H)$ with $|U| = m \leq \varepsilon n$ and m is a prime number. Then there exists a $C_\ell^{(k)}$ -decomposable J_0 such that $|J_0| = \ell m$, $J_0[U] = \emptyset$, J_0 has a balanced tour-trail decomposition \mathcal{T}_0 satisfying $V(D(\mathcal{T}_0)) \subseteq U$ and, for all $1 \leq i < k/2$, $A_i(\mathcal{T}_0)$ is a strongly connected multidigraph which spans U and $|A_i(\mathcal{T}_0)| = 2m$.*

Proof. Let u_1, \dots, u_m be an enumeration of U . Consider $j \in [m]$. We apply Lemma 6.1 to obtain a copy C_j of $C_\ell^{(k)}$ with $V(C_j) = u_{j+1} \cdots u_{j+k-1} w_{j,k} \cdots w_{j,\ell}$, where $w_{j,j'}$ are new vertices. For its trail decomposition \mathcal{T}_j we consider C_j to be a trail

$$u_{j+1} \cdots u_{j+k-1} w_{j,k} \cdots w_{j,\ell} u_{j+1} \cdots u_{j+k-1}.$$

Then $u_{j+i} u_{j+k-i}, u_{j+k-i} u_{j+i} \in A_i(\mathcal{T}_j)$ for all $1 \leq i < k/2$. Let $J_0 = \bigcup_{j \in [m]} C_j$ and $\mathcal{T}_0 = \bigcup_{j \in [m]} \mathcal{T}_j$ (without simplification). Note that $|J_0| = \ell m$ as each C_j has ℓ edges. Note that

$$|A_i(\mathcal{T}_0)| = \sum_{j \in [m]} |A_i(\mathcal{T}_j)| = \sum_{j \in [m]} |D(\mathcal{T}_j)| = 2m.$$

Recall that m is prime, so $A_i(\mathcal{T}_0)$ is connected and spans U for all $1 \leq i < k/2$. \blacksquare

We are now ready to prove Lemma 7.6.

Proof of Lemma 7.6. Apply Lemma 7.8 with $U = V(G)$ and $H = H \setminus \{z_1, \dots, z_{k-1}\}$, we obtain a $C_\ell^{(k)}$ -decomposable graph $J_0 \subseteq H - G$ such that

$$(7.4) \quad |J_0| = \ell m,$$

and J_0 has a balanced tour-trail decomposition \mathcal{T}_0 such that, for all $1 \leq i < k/2$, $A_i(\mathcal{T}_0)$ is a connected multidigraph which spans $V(G)$.

Let $G_0^* = G \cup J_0$ and $\mathcal{T}_0^* = \mathcal{T} \cup \mathcal{T}_0$, considering all trails, without doing any further simplification even if it is possible to do so. For all $1 \leq i < k/2$, $A_i = A_i(\mathcal{T}_0^*)$. Thus A_i is a connected multidigraph spanning $V(G)$. Observe that since \mathcal{T}_0^* is balanced, then A_i is Eulerian. Let s be such that $|D(\mathcal{T}_0^*)| = 2s$. Note that

$$(7.5) \quad 2s = |A_i| = |D(\mathcal{T}_0^*)| = 2|G| + |A_i(\mathcal{T}_0)| \leq 2m/k + 2m \leq 3m.$$

Claim 7.9. *There exist edge-disjoint k -graphs $J_0, \dots, J_{\lfloor k/2 \rfloor}$ in $H - G$ such that*

- (i) *each J_i is $C_\ell^{(k)}$ -decomposable and $|J_i| \leq 2 \cdot 3^i k \ell s$,*
- (ii) *$G \cup J_0 \cup \bigcup_{j \in [i]} J_j$ has a balanced tour-trail decomposition \mathcal{T}_i^* , and*
- (iii) *\mathcal{T}_i^* is an i -convert of \mathcal{T}_{i-1}^* .*

We first show that the claim implies the lemma. Set $J^* = J_0 \cup \bigcup_{j \in \llbracket k/2 \rrbracket} J_j$ and set $\mathcal{T}^* = \mathcal{T}_{\llbracket k/2 \rrbracket}^*$. Clearly, J^* is $C_\ell^{(k)}$ -decomposable since each J_j is. Note that

$$|J^*| \leq |J_0| + \sum_{i \in \llbracket k/2 \rrbracket} |J_i| \stackrel{(7.4),(i)}{\leq} \ell m + 2k\ell s \sum_{i \in \llbracket k/2 \rrbracket} 3^i \stackrel{(7.5)}{\leq} 3^k k\ell m.$$

Since \mathcal{T}^* is balanced by (ii), (iii) implies that each $v_1 \cdots v_{k-1} \in D(\mathcal{T}^*)$ satisfies

$$v_1 \cdots v_{k-1} \in \{z_1, z_{k-1}\} \times \{z_2, z_{k-2}\} \times \cdots \times \{z_{k-1}, z_1\}.$$

Hence, for all $i \in [k-1]$ and $v \in V(H)$, $p_{\mathcal{T}^*,i}(v) = 0$ unless $v \in \{z_i, z_{k-i}\}$ as desired. Also $|D(\mathcal{T}^*)| = |D(\mathcal{T})| = 2s \leq 3m$ by (iii) and (7.5). Therefore to complete the proof, it remains to prove Claim 7.9.

Proof of claim. Suppose that we have already found J_0, \dots, J_{i-1} and we now construct J_i as follows.

Case 1: $i = k/2$. We first prove the case $i = k/2$ as it is simpler and illustrates some of the key ideas. If we are in this case, then k is even and, by Proposition 7.2, we have $p_{\mathcal{T}_{k/2-1}, k/2}(v) \equiv 0 \pmod{2}$ for all $v \in V(H)$. Also $|D(\mathcal{T}_{k/2-1}^*)| = |\mathcal{T}_0^*| = 2s$. Take an arbitrary enumeration of $D(\mathcal{T}_{k/2-1}^*)$ into $\mathbf{b}_1, \dots, \mathbf{b}_{2s}$ such that $b_{2j-1, k/2} = b_{2j, k/2}$ for all $j \in [s]$, where $\mathbf{b}_j = b_{j,1} \cdots b_{j,k-1}$ (here we use the fact that $p_{\mathcal{T}_{k/2-1}, k/2}(v)$ is even). For every $j \in [s]$, let

$$b_j^* = b_{2j-1, k/2} = b_{2j, k/2}.$$

Apply Lemma 6.6 to obtain a swapper gadget $T_{k/2}^j = T_{k/2}(\mathbf{b}_{2j-1}, \mathbf{b}_{2j}^{-1}, z_{k/2}, b_j^*)$ with a tour-trail decomposition \mathcal{T}^j such that

$$D(\mathcal{T}^j) = \bar{T}_{k/2}(\mathbf{b}_{2j-1}, \mathbf{b}_{2j}^{-1}, z_{k/2}, b_j^*) = \{\mathbf{b}_{2j-1}^{-1}, \mathbf{b}_{2j}^{-1}, \zeta_{k/2}(\mathbf{b}_{2j-1}), \zeta_{k/2}(\mathbf{b}_{2j})\}.$$

We may further assume that these T^j are edge-disjoint.

Let $J_{k/2}$ be the union of these swapper gadgets and $\mathcal{T}_{k/2}$ be the union of their tour-trail decompositions, together with $\mathcal{T}_{k/2-1}$. Note that $|J_{k/2}| \leq 3^{k/2} \ell k s$, and since

$$\begin{aligned} D(\mathcal{T}_{k/2}^*) &= D(\mathcal{T}_{k/2-1}^*) \cup \bigcup_{j \in [s]} \{\mathbf{b}_{2j-1}^{-1}, \mathbf{b}_{2j}^{-1}, \zeta_{k/2}(\mathbf{b}_{2j-1}), \zeta_{k/2}(\mathbf{b}_{2j})\} \\ &= \{\mathbf{b}_j, \mathbf{b}_j^{-1}, \zeta_{k/2}(\mathbf{b}_j) : j \in [2s]\} = \{\zeta_{k/2}(\mathbf{b}_j) : j \in [2s]\}, \end{aligned}$$

we deduce $\mathcal{T}_{k/2}$ is a $(k/2)$ -convert of $\mathcal{T}_{k/2-1}$, as required.

Case 2: $i < k/2$. By (iii) and Proposition 7.7, we deduce that $A_i(\mathcal{T}_{i-1}) = A_i$ and so, it is an Eulerian multidigraph. Hence, there exists an enumeration of $D(\mathcal{T}_i)$ as $\mathbf{b}_1, \dots, \mathbf{b}_{2s}$, which corresponds to an Eulerian tour in A_i . Let $\mathbf{b}_j = b_{j,1} \cdots b_{j,k-1}$, so we have $b_{j,k-i} = b_{j+1,i}$ for all $j \in [2s]$.

We now replace the $(k-i)$ th vertex in each b_{2j-1} with z_i and the i th vertex in each b_{2j} with z_i , as follows. For every $j \in [s]$ let

$$b_j^* = b_{2j-1, k-i} = b_{2j, i}.$$

Apply Lemma 6.6 to obtain a swapper gadget $T^j = T_i(\mathbf{b}_{2j}, \mathbf{b}_{2j-1}^{-1}, z_i, b_j^*)$ with a tour-trail \mathcal{T}^j such that

$$D(\mathcal{T}^j) = \bar{T}_i(\mathbf{b}_{2j}, \mathbf{b}_{2j-1}^{-1}, z_i, b_j^*) = \{\mathbf{b}_{2j-1}^{-1}, \mathbf{b}_{2j}^{-1}, r_{k-i}(\mathbf{b}_{2j-1}, z_i), r_i(\mathbf{b}_{2j}, z_i)\}.$$

Recall that $r_i(\mathbf{b}_{2j}, z_i)$ and $r_{k-i}(\mathbf{b}_{2j-1}, z_i)$ correspond to the tuples \mathbf{b}_{2j} and \mathbf{b}_{2j-1} replacing b_j^* with z_i (see definition at the beginning of Section 6). Note that

$$D(\mathcal{T}_i \cup \bigcup_{j \in [s]} \mathcal{T}^j) = \bigcup_{j \in [2s]} \{\mathbf{b}_{2j-1}, \mathbf{b}_{2j}, \mathbf{b}_{2j-1}^{-1}, \mathbf{b}_{2j}^{-1}, r_{k-i}(\mathbf{b}_{2j-1}, z_i), r_i(\mathbf{b}_{2j}, z_i)\}$$

$$= \bigcup_{j \in [s]} \{r_{k-i}(\mathbf{b}_{2j-1}, z_i), r_i(\mathbf{b}_{2j}, z_i)\}.$$

Equivalently, $D(\mathcal{T}_i \cup \bigcup_{j \in [2s]} \mathcal{T}^j)$ is obtained from $D(\mathcal{T}_i)$ by corresponds to replacing the $(k-i)$ th vertex in \mathbf{b}_{2j-1} and i th vertex in \mathbf{b}_{2j} with z_i .

By considering the pairs of tuples $r_i(\mathbf{b}_{2j}, z_i), r_{k-i}(\mathbf{b}_{2j+1}, z_i)$, a similar argument implies that we can replace the i th vertex in $r_{k-i}(\mathbf{b}_{2j+1}, z_i)$ and $(k-i)$ th vertex in $r_i(\mathbf{b}_{2j}, z_i)$ with z_{k-i} .

Let J_i be the union of these swapper gadgets and \mathcal{T}_i be the union of \mathcal{T}_{i-1} and the corresponding tour-trail decomposition. Notice that \mathcal{T}_i is an i -convert of \mathcal{T}_{i-1} and that $|J_i| \leq 3^j \ell k \cdot 2s$. This finishes the proof of Claim 7.9. \square

As discussed, this finishes the proof of the lemma. \blacksquare

7.4. Untangle the last arcs. Observe that Lemma 7.6 in the previous subsection we found a trail-tour decomposition in which all arcs lie in a small set of $k-1$ vertices z_1, \dots, z_{k-1} . Here we show how to ‘untangle’ those arcs in such a way that all ‘cancel’ each other. After this cancelling, the trails from the tour-trail decomposition are removed and we obtain a tour decomposition.

Lemma 7.10. *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k$ with $k \geq 3$ and $\ell \geq k^2 - k + 1$. Let H be a k -graph on n vertices with $\delta^{(3)}(H) \geq \varepsilon n$. Let G be a k -graph with $V(G) \subseteq V(H)$ and $|V(G)| \leq \varepsilon n$. Let $z_1, \dots, z_{k-1} \in V(G)$ be distinct vertices. Suppose that G has a balanced tour-trail decomposition \mathcal{T}_1 such that $|D(\mathcal{T}_1)| \leq 5m$ and, for all $i \in [k-1]$ and $v \in V(H)$, $p_{\mathcal{T}_1, i}(v) = 0$ unless $v \in \{z_i, z_{k-i}\}$. Then $H - G$ contains a $C_\ell^{(k)}$ -decomposable subgraph J such that $|J| \leq k^3 \ell |D(\mathcal{T}_1)|$, $J[V(G)] = \emptyset$, and $G \cup J$ has a tour decomposition.*

Proof. We simplify \mathcal{T}_1 as much as possible. Let $|D(\mathcal{T}_1)| = 2s$. Since \mathcal{T}_1 is balanced, we have, for all $i \in [k-1]$,

$$(7.6) \quad p_{\mathcal{T}_1, i}(z_i) = p_{\mathcal{T}_1, i}(z_{k-i}) = \begin{cases} s & \text{if } i \neq k/2, \\ 2s & \text{if } i = k/2. \end{cases}$$

We now colour $\mathbf{b} \in D(\mathcal{T}_1)$ red if \mathbf{b} starts at z_1 (i.e. the first vertex of \mathbf{b} is z_1), and blue otherwise. So there are s red $(k-1)$ -tuples and s blue $(k-1)$ -tuples in $D(\mathcal{T}_1)$. Ideally, we would like to transform all red $(k-1)$ -tuples to $z_1 \cdots z_{k-1}$ and all blue $(k-1)$ -tuples to $z_{k-1} \cdots z_1$ (so that they would cancel out). A $(k-1)$ -tuple is i -bad if z_i is at the ‘wrong place’. More precisely, an i -bad red $(k-1)$ -tuple (and an i -bad blue $(k-1)$ -tuple) will be of form $v_1 \cdots v_{i-1} z_{k-i} v_{i+1} \cdots v_{k-i-1} z_i v_{k-i+1} \cdots v_{k-1}$ (and $v_1 \cdots v_{i-1} z_i v_{i+1} \cdots v_{k-i-1} z_{k-i} v_{k-i+1} \cdots v_{k-1}$, respectively), where $\{v_j, v_{k-j}\} = \{z_j, z_{k-j}\}$ for all $j \in [k-1] \setminus \{i\}$. Note that no $(k-1)$ -tuple is 1-bad, $(k-1)$ -bad or $k/2$ -bad. If a $(k-1)$ -tuple is i -bad, then it is also $(k-i)$ -bad. By (7.6), the number of red i -bad paths equals to the number of blue i -bad paths.

We claim that there exist edge-disjoint k -graphs $J_2, \dots, J_{\lfloor k/2 \rfloor - 1}$ in $G - H$ such that, for $2 \leq i < k/2$,

- (i) each J_i is $C_\ell^{(k)}$ -decomposable and $|J_i| \leq 8i \ell k s$,
- (ii) $G \cup \bigcup_{j \in [2, i]} J_j$ has a balanced tour-trail decomposition \mathcal{T}_i ,
- (iii) for all $j \in [k-1]$ and $v \in V(H)$, $p_{\mathcal{T}_i, j}(v) = 0$ unless $v \in \{z_j, z_{k-j}\}$,
- (iv) $|D(\mathcal{T}_i)| = |D(\mathcal{T}_{i-1})|$, and
- (v) $D(\mathcal{T}_i)$ contains no j -bad $(k-1)$ -tuple for all $j \in [i]$.

Suppose that, for some $2 \leq i < k/2 - 1$, we have constructed J_2, \dots, J_{i-1} . We describe the construction of J_i as follows.

Let $\mathbf{a}_1, \dots, \mathbf{a}_t$ be the i -bad red $(k-1)$ -tuples in $D(\mathcal{T}_{i-1})$ and $\mathbf{b}_1, \dots, \mathbf{b}_t$ be the i -bad blue $(k-1)$ -tuples in $D(\mathcal{T}_{i-1})$. Consider any $j \in [t]$. Let

$$\mathbf{a}_j = a_{j,1} \cdots a_{j,i-1} z_{k-i} a_{j,i+1} \cdots a_{j,k-i-1} z_i a_{j,k-i+1} \cdots a_{j,k-1}$$

$$\mathbf{b}_j = b_{j,1} \cdots b_{j,i-1} z_i b_{j,i+1} \cdots b_{j,k-i-1} z_{k-i} b_{j,k-i+1} \cdots b_{j,k-1}.$$

We now mimic the argument in the proof of Lemma 7.6 to swap z_i and z_{k-i} in \mathbf{a}_j 's and \mathbf{b}_j 's. However, we are unable to construct swapper gadgets $T_i(\mathbf{a}_j, \mathbf{b}_j^{-1}, z_i, z_{k-i})$ as both \mathbf{a}_j and \mathbf{b}_j contain both z_i and z_{k-i} . To overcome this problem, we first replace z_i with a new vertex w (so \mathbf{a}_j and \mathbf{b}_j are now free of z_i). After this is done, then we can replace z_{k-i} with z_i , and finally we replace w with z_{k-i} . We now formalise the proof as follows.

Let $w \in V(H) \setminus \{z_{i'} : i' \in [k-1]\}$ be a new vertex. Apply Lemma 6.6 to obtain three edge-disjoint swapper gadgets

$$T_i(\mathbf{a}_j, \mathbf{b}_j^{-1}, w, z_{k-i}), \quad T_i(r_i(\mathbf{a}_j, w)^{-1}, r_{k-i}(\mathbf{b}_j, w), z_i, z_{k-i}) \quad \text{and} \\ T_i(r_{k-i}(\mathbf{a}_j, z_{k-i}), r_i(\mathbf{b}_j, z_{k-i})^{-1}, z_i, w)$$

Let T^j be its union and \mathcal{T}^j be the union of their tour-trail decompositions. It is not hard to check that after cancellation we obtain

$$D(\mathcal{T}^j) = \{\mathbf{a}_j^{-1}, \mathbf{b}_j^{-1}, \zeta_i(\mathbf{a}_j), \bar{\zeta}_i(\mathbf{b}_j)\}.$$

Note that $\zeta_i(\mathbf{a}_j)$ and $\bar{\zeta}_i(\mathbf{b}_j)$ are not i -bad.

Let $J_i = \bigcup_{j \in [t]} T^j$ and let $\mathcal{T}_i = \mathcal{T}_{i-1} \cup \bigcup_{j \in [t]} \mathcal{T}^j$ be the corresponding tour-trail decomposition of $G \cup \bigcup_{2 \leq j \leq i} J_j$. This finishes the construction of J_i .

Now we have constructed J_i for all $2 \leq i < k/2$. We set $J = \bigcup_{2 \leq i < k/2} J_i$, so $|J| \leq 2k^3 \ell s = k^3 \ell |D(\mathcal{T}_1)|$. Note that $\mathcal{T}_{\lfloor k/2 \rfloor - 1}$ is a balanced tour-trail decomposition of $G \cup J$ without any bad $(k-1)$ -tuple. Therefore, after cancellation, $D(\mathcal{T}_{\lfloor k/2 \rfloor - 1})$ is empty, implying that $\mathcal{T}_{\lfloor k/2 \rfloor - 1}$ is a tour decomposition. \blacksquare

7.5. Proof of Lemma 7.1. We now put the pieces together to prove Lemma 7.1.

Proof of Lemma 7.1. Apply Lemma 7.3 to obtain a $C_\ell^{(k)}$ -decomposable J_1 in $H - G$ such that $|G \cup J_1| \leq \ell m^{k+1}$ and $G \cup J$ has a balanced tour-trail decomposition \mathcal{T}_1 .

Let m_1 be a prime between $k\ell m^{k+1}$ and $2k\ell m^{k+1}$ (this exists by Bertrand's postulate). Add isolated vertices to $G \cup J_1$ to obtain a subgraph G_1 of H such that

$$|V(G_1)| = m_1 \text{ and } |G_1| \leq m_1/k.$$

Let $z_1, \dots, z_{k-1} \in V(H) \setminus V(G_1)$. Apply Lemma 7.6 (with G_1, \mathcal{T}_1 playing the rôles of G, \mathcal{T}) to obtain a $C_\ell^{(k)}$ -decomposable J_2 in $H - G_1$ such that

$$|J_2| \leq 3^k k \ell m_1$$

and $G_2 = G_1 \cup J_2$ has a balanced tour-trail decomposition \mathcal{T}_2 satisfying, for all $i \in [k-1]$ and $v \in V(H)$, $p_{\mathcal{T}_2, i}(v) = 0$ unless $v \in \{z_i, z_{k-i}\}$ and $|D(\mathcal{T}_2)| \leq 3m_1$.

Apply Lemma 7.10 (with G_2, \mathcal{T}_2 playing the rôles of G, \mathcal{T}_1) to acquire a $C_\ell^{(k)}$ -decomposable subgraph J_3 in $H - G_2$ such that

$$|J_3| \leq k^3 \ell |D(\mathcal{T}_2)| \leq 3k^3 \ell m_1$$

and $G_2 \cup J_3$ has a tour decomposition. We set $J = J_1 \cup J_2 \cup J_3$ and so $G \cup J$ has a tour decomposition and

$$|G \cup J| \leq m_1 + 3^k k \ell m_1 + 3k^3 \ell m_1 \leq 3^{k+2} k^4 \ell^2 m^{k+1}.$$

The 'moreover' statement can be obtained by constructing a homomorphic copy of J in $H - G - J$. \blacksquare

8. TRANSFORMERS III: PROOF OF LEMMA 4.3

Finally, in this section we use the machinery of transformers and tour-trail decompositions to find cycle absorbers and prove Lemma 4.3.

Proof of Lemma 4.3. Let $W \subseteq V(H)$ be of size at $m \leq m_0$. By Lemma 6.1, for any two ordered $(k-1)$ -tuples $v_1 \cdots v_k$ and $v_{\ell-k+1} \cdots v_\ell$ in $V(H) \setminus W$, $H \setminus W$ contains a tight walk $v_1 \cdots v_\ell$ on ℓ vertices and no repeated vertices outside of $v_1 \cdots v_k$ and $v_{\ell-k+1} \cdots v_\ell$. In particular, there is an ℓ -cycle in $H \setminus W$ covering any arbitrary k -tuple v_1, \dots, v_k in $H \setminus W$. Hence, by Lemma 3.3, it suffices to show that H is $(C_\ell^{(k)}, m_0, m_0, \eta)$ -transforming for some increasing function $\eta : \mathbb{N} \rightarrow \mathbb{N}$ which satisfies $\eta(x) \geq x$. Indeed, this will imply that H is $(C_\ell^{(k)}, m_0, m_0, \eta')$ -absorbable for some increasing function $\eta' : \mathbb{N} \rightarrow \mathbb{N}$ such that $\eta'(x) \geq x$, as desired.

Define $\eta : \mathbb{N} \rightarrow \mathbb{N}$ by $\eta(x) = 3^{k+3}k^5\ell^3x^{k+1}$. We will show that H is $(C_\ell^{(k)}, m_0, m_0, \eta)$ -transforming. To do so, let G_1 and G_2 be two vertex-disjoint $C_\ell^{(k)}$ -divisible k -graphs with $V(G_1), V(G_2) \subseteq V(H)$; suppose that $|V(G_1)|, |V(G_2)| \leq m_0$ and that there is an edge-bijective homomorphism from G_1 to G_2 . Let $W \subseteq V(H) \setminus V(G_1 \cup G_2)$ with $|W| \leq m_0$. Let $m = \max\{|V(G_1)|, |V(G_2)|\}$. Let $H' = H \setminus W$. It is enough to show that H' contains a $(G_1, G_2; C_\ell^{(k)})$ -transformer of order at most $\eta(m)$. This will be our task from now on.

Apply Lemma 7.1 to obtain vertex-disjoint subgraphs J_1 and J_2 of $H' - G_1 - G_2$ such that

- (a₁) $G_1 \cup J_1$ and $G_2 \cup J_2$ have tour decompositions,
- (a₂) J_1 and J_2 are $C_\ell^{(k)}$ -decomposable,
- (a₃) $J_1[V(G_1 \cup G_2)]$ and $J_2[V(G_1 \cup G_2)]$ are empty,
- (a₄) $|G_1 \cup J_1|, |G_2 \cup J_2| \leq 3^{k+2}k^4\ell^2m^{k+1}$, and
- (a₅) there is an edge-bijective homomorphism ϕ from $G_1 \cup J_1$ to $G_2 \cup J_2$.

Let $G'_j = G_j \cup J_j$ for $j \in [2]$. We now claim that there exists a $(G'_1, G'_2; C_\ell^{(k)})$ -transformer T^* with $|T_i| = (\ell-1)|G'_1|$. Indeed, let $\{A_i : i \in [s]\}$ be a tour decomposition of G'_1 , and recall that ϕ is an edge-bijective homomorphism from G'_1 to G'_2 . Therefore, $\{\phi(A_i) : i \in [s]\}$ is a tour decomposition of G'_2 . Now, suppose that for some $i \in [s]$, we have already constructed edge-disjoint T_1, \dots, T_{i-1} in $H' - G'_1 - G'_2$ such that, for $i' \in [i-1]$

- (b₁) $T_{i'}$ is an $(A_{i'}, \phi(A_{i'}); C_\ell^{(k)})$ -transformer,
- (b₂) $|T_{i'}| = (\ell-1)|A_{i'}|$, and
- (b₃) $T_{i'}[V(G'_1)] \cup T_{i'}[V(G'_2)] = \emptyset$.

We construct T_i as follows. Let $H^* = H' - G'_1 - G'_2 - \bigcup_{i' \in [i-1]} T_{i'}$. Let $A_i = v_1 \cdots v_t$. Let $w_j = \phi(v_j)$ for all $j \in [t]$, so $\phi(A_i) = w_1 \cdots w_t$. Note that $\ell - k^2 + k - 1 \geq k^2 - k$ by the choice of ℓ . Therefore, by Lemma 6.1, H^* contains tight paths $P_1, \dots, P_s, Q_1, \dots, Q_s$ such that, for each $j \in [t]$,

- (c₁) P_j is a tight path of length $k^2 - k$ from $v_{j+1}v_{j+2} \cdots v_{j+k-1}$ to $w_jw_{j+1} \cdots w_{j+k-2}$,
- (c₂) Q_j is a tight path of length $\ell - k^2 + k - 1$ starting from $w_jw_{j+1} \cdots w_{j+k-2}$ and ending in $v_jv_{j+1} \cdots v_{j+k-2}$, and
- (c₃) $V(P_j) \setminus V(D(P_j))$ and $V(Q_j) \setminus V(D(Q_j))$ are new vertices.

Each of $v_jv_{j+1} \cdots v_{j+k-1} \cup P_j \cup Q_j$ and $w_jw_{j+1} \cdots w_{j+k-1} \cup P_j \cup Q_{j+1}$ forms a $C_\ell^{(k)}$. Hence, we are done by letting $T_i = \bigcup_{j \in [s]} P_j \cup Q_j$. This finishes the construction of T_i . Thus $T^* = \bigcup_{i \in [s]} T_i$ is the desired $(G'_1, G'_2; C_\ell^{(k)})$ -transformer.

We finish by defining $T = J_1 \cup J_2 \cup T^*$, so

$$|V(T)| \leq k|T| \leq k(\ell+1)|G'_1| \stackrel{(a_4)}{\leq} 3^{k+3}k^5\ell^3m^{k+1} = \eta(m).$$

Together with (a₂) and (a₃), we deduce that $T[V(G_1)]$ is empty and

$$G_1 \cup T = ((G_1 \cup J_1) \cup T^*) \cup J_2 = (G'_1 \cup T^*) \cup J_2$$

is $C_\ell^{(k)}$ -decomposable, and similarly $G_2 \cup T$ is $C_\ell^{(k)}$ -decomposable. Therefore, T is a $(G_1, G_2; C_\ell^{(k)})$ -transformer. \blacksquare

9. COVER-DOWN LEMMA

In this section we prove Lemma 4.2 which is the main step in the iterative part of iterative absorption. We prove this lemma by induction on k , and when dealing with k -uniform hypergraphs we will require results on path decompositions for $k - 1$. To organise our arguments, we define the following two statements for each $k \geq 3$. The first statement corresponds precisely to the Cover-down lemma for k -graphs, while the second one concerns decompositions of k -graphs into paths.

(\odot_k) For every $\alpha > 0$, there is an $\ell_0 \in \mathbb{N}$ such that for every $\mu > 0$ and every $n, \ell \in \mathbb{N}$ with $\ell \geq \ell_0$ and $1/n \ll \mu, \alpha$ the following holds. Let H be a k -graph on n vertices and $U \subseteq V(H)$ with $|U| = \lfloor \alpha n \rfloor$ such that

$$(CD_1) \quad \delta^{(2)}(H) \geq 2\alpha n,$$

$$(CD_2) \quad \delta^{(2)}(H, U) \geq \alpha|U| \text{ and}$$

$$(CD_3) \quad \deg_H(x) \text{ is divisible by } k \text{ for each } x \in V(H) \setminus U.$$

Then H contains a $C_\ell^{(k)}$ -decomposable subgraph $F \subseteq H$ such that $H - H[U] \subseteq F$ and $\Delta_{k-1}(F[U]) \leq \mu n$.

(\curvearrowright_k) For every $\ell \geq k$ and for every $\alpha > 0$ there is an n_0 such that the following holds. Every k -graph H on $n \geq n_0$ vertices with $\delta^{(2)}(H) \geq \alpha n$ and $|H| \equiv 0 \pmod{\ell}$ contains a $P_\ell^{(k)}$ -decomposition.

Thus, Lemma 4.2 can be synthetically stated as follows.

Lemma 4.2 (Cover-down lemma (reprise)). (\odot_k) holds for every $k \geq 3$.

We show Lemma 4.2 through an induction on k , in which (\curvearrowright_k) is helpful to enable the induction step.

Lemma 9.1. For each $k \geq 3$, if (\curvearrowright_{k-1}) holds, then (\odot_k) holds.

Lemma 9.2. For each $k \geq 3$, if (\odot_k) holds, then (\curvearrowright_k) holds.

Assuming the validity of these two last lemmas, Lemma 4.2 follows easily, if we are provided with a base case. For this, we use the following result by Botler, Mota, Oshiro and Wakabayashi [4] on path decomposition in graphs, which immediately yields (\curvearrowright_2).

Theorem 9.3 ([4]). For each $\ell \geq 1$, there exists k_ℓ such that each k_ℓ -edge-connected graph whose number of edges is divisible by ℓ has a $P_\ell^{(2)}$ -decomposition.

The proof of Lemma 9.2 is given in the next subsection. The proof of Lemma 9.1 will require more effort and is given in Subsection 9.4, after some previous necessary results.

9.1. Path decompositions: proof of Lemma 9.2 and Theorem 1.4. To see that the bound $\delta_{P_\ell^{(k)}} \geq 1/2$ holds, consider the following example. Take the union of $K_{\lfloor n/2 \rfloor}^{(k)}$ and $K_{\lfloor n/2 \rfloor}^{(k)}$ on vertex sets A and B , respectively. Delete a few edges if necessary so that resulting k -graph H satisfies $|H| \equiv 0 \pmod{\ell}$ but $|H[A]| \not\equiv 0 \pmod{\ell}$. Then H is not $P_\ell^{(k)}$ -decomposable and $\delta(H) \geq (1/2 - o(1))n$. On the other hand, note that Lemma 9.2 and Lemma 9 imply Theorem 1.4.

The proof of Lemma 9.2 follows essentially the same strategy we use to prove Theorem 1.3 in Section 4. The Vortex lemma is the same and for the Cover-down lemma we may use (\odot_k), which is assumed to hold as a hypothesis. To see this, it is enough to notice that for every sufficiently large ℓ' divisible by ℓ , a $C_{\ell'}^{(k)}$ -decomposable subgraph is $P_\ell^{(k)}$ -decomposable as well. Hence, the only new ingredient needed is the following Absorber lemma for paths.

Lemma 9.4 (Absorber lemma for paths). *Let $1/n \ll \varepsilon \ll 1/\ell, 1/k, 1/m_0$ with $k \geq 3$. Let H be a k -graph on n vertices with $\delta^{(2)}(H) \geq \varepsilon n$. Then H is $(P_\ell^{(k)}, m_0, m_0, \eta')$ -absorbing for some increasing function $\eta' : \mathbb{N} \rightarrow \mathbb{N}$ satisfying $\eta'(x) \geq x$ and independent of ε and n .*

Proof. Pick an arbitrary edge $e \in P_\ell^{(k)}$ and let \mathbf{a} be any k -tuple in $V(H)$. Since the inequality $\delta^{(2)}(H) \geq \varepsilon n$ holds, it is easy to see for every $W \subseteq V(H) \setminus \{v_1, \dots, v_k\}$ of size at most m_0 there is a copy of $P_\ell^{(k)}$ in $(H \cup \{\mathbf{a}\}) \setminus W$ in which \mathbf{a} plays the rôle of e . We do this by simply extending a path (maybe in both directions) starting with \mathbf{a} , which we can do simply because of $\delta(H) \geq \delta^{(2)}(H) \geq \varepsilon n$. This enables us to use Lemma 3.3, and hence, it is enough to prove that H is $(P_\ell^{(k)}, m_0, m_0, \eta)$ -transforming for some increasing function $\eta : \mathbb{N} \rightarrow \mathbb{N}$.

Let ℓ_0 be the smallest number larger than $k^2 - k$ which is divisible by ℓ and define $\eta(x) = \ell_0 x^2$. Let $G, G' \subseteq H$ be vertex-disjoint $P_\ell^{(k)}$ -divisible subgraphs such that there is an edge-bijective homomorphism ϕ from G to G' . Also, let $W \subseteq V(H) \setminus V(G \cup G')$ and suppose $|V(G)|, |V(G')| \leq m_0$ and $|W| \leq m_0 - \eta(|V(G)|)$. For every edge $e \in G$ apply Lemma 6.1 to find a path $P_e \subseteq H \setminus W$ between e and $\phi(e)$ with precisely $\ell_0 + 1$ edges. Since ℓ_0 is divisible by ℓ , $T = \bigcup_{e \in G} P_e$ is a $(G, G'; P_\ell^{(k)})$ -transformer of size at most $\ell_0 e(G) \leq \eta(\max\{|V(G)|, |V(G')|\})$. ■

We omit further details of proof of Lemma 9.2 and reference the reader to the proof of Theorem 1.3.

9.2. Well-behaved approximate cycle decompositions. Given a k -graph H such that $\delta^{(2)}(H) \geq \alpha n$, we find a $C_\ell^{(k)}$ -packing \mathcal{C} that covers almost all edges of H and such that the leftover is not too concentrated in a $(k-1)$ -tuple. Here, a $C_\ell^{(k)}$ -packing is a set of edge-disjoint copies of $C_\ell^{(k)}$. More precisely, we have the following lemma.

Lemma 9.5 (Well-behaved cycle decompositions). *Given $k \in \mathbb{N}$ and $\alpha \geq 0$ there is an $\ell_0 \in \mathbb{N}$ such that for every $\gamma > 0$ and $\ell, n \in \mathbb{N}$ with $\ell \geq \ell_0$ and $1/n \ll \gamma, \alpha, 1/\ell$ the following holds. Let H be a k -graph on n vertices with $\delta^{(2)}(H) \geq \alpha n$. Then H has a $C_\ell^{(k)}$ -packing \mathcal{C} such that $\Delta_{k-1}(H - \bigcup \mathcal{C}) \leq \gamma n$.*

The case $k = 3$ is proven by the last two authors in [14] and here we follow the same lines. Given a k -graph H and an edge $e \in H$, recall that $\mathcal{C}_\ell(H)$ and $\mathcal{C}_\ell(H, e)$ are the family of all ℓ -cycles in H and those containing e . The proof of Lemma 9.5 rests in a result by Joos and Kühn [11] about fractional C_ℓ^k -decompositions.

Theorem 9.6 (Joos and Kühn [11]). *Given $k \in \mathbb{N}$ and $\alpha, \mu \geq 0$ there is an $\ell_0 \in \mathbb{N}$ such that for every $\ell, n \in \mathbb{N}$ with $\ell \geq \ell_0$ and $1/n \ll \alpha, 1/\ell$ the following holds. Let H be a k -graph on n vertices with $\delta^{(2)}(H) \geq \alpha n$. Then there is a fractional $C_\ell^{(k)}$ -decomposition ω of H with*

$$(1 - \mu) \frac{2|H|}{\Delta_{k-1}(H)^\ell} \leq \omega(C) \leq (1 + \mu) \frac{2|H|}{\delta_{k-1}(H)^\ell}$$

for all cycles $C \in \mathcal{C}_\ell(H)$.

Additionally, we need the following nibble-type matching theorem. The statement of the theorem is technical, but in our context the conditions are easy to check. Consider the following parameter $g(H) = \Delta_1(H)/\Delta_2(H)$ for every k -graph H .

Theorem 9.7 (Alon and Yuster [1]). *For every $\gamma > 0$, there is a $\xi > 0$ such that for every sufficiently large n the following holds. Let H be a k -graph on n vertices and let $U_1, \dots, U_t \subseteq V(H)$ be subsets of vertices with $\log t \leq g(H)^{1/(3k-3)}$ and such that $|U_i| \geq 5g(H)^{1/(3k-3)} \log(g(H)t)$ for every $i \in [t]$. Suppose that*

- (a) $\delta_1(H) \geq (1 - \xi)\Delta_1(H)$ and
- (b) $\Delta_1(H) \geq (\log n)^7 \Delta_2(H)$.

Then H contains a matching such that at most $\gamma|U_i|$ vertices are uncovered in each U_i .

Lemma 9.5 follows by an straightforward application of Theorems 9.6 and 9.7.

Proof of Lemma 9.5. Given $k \in \mathbb{N}$ and $\alpha > 0$ fix any $\mu, \xi < 1/3$ and take ℓ_0 given by Theorem 9.6. Let $\ell \geq \ell_0$, $\gamma > 0$ and let n be sufficiently large for an application of Theorems 9.6 and 9.7.

First, we apply Theorem 9.6 to obtain a fractional $C_\ell^{(k)}$ -decomposition ω of H satisfying

$$(9.1) \quad \omega(C) \leq (1 + \mu) \frac{2|H|}{\delta_{k-1}(H)^\ell} \leq \frac{3n^k}{\delta^{(2)}(H)^\ell} \leq \frac{3}{\alpha^\ell n^{\ell-k}},$$

for all cycles $C \in \mathcal{C}_\ell(H)$.

Then, consider the auxiliary ℓ -graph F with vertex set $E(H)$ and an edge in F for each cycle in $\mathcal{C}_\ell(H)$ corresponding to its set of ℓ edges in H . Define a random subgraph $F' \subseteq F$ by keeping each edge C with probability $p_C = n^{1/2}\omega(C) \leq 1$ by (9.1).

For every edge $e \in H$ we have $\mathbb{E}[d_{F'}(e)] = n^{1/2} \sum_{C \in \mathcal{C}_\ell(H, e)} \omega(C) = n^{1/2}$. Moreover, since two distinct edges $e, f \in E(H)$ can participate together in at most $O(n^{\ell-(k+1)})$ many $C_\ell^{(k)}$ in H , (9.1) implies that the expected 2-degree is bounded by $\mathbb{E}[d_{F'}(e, f)] = O(n^{-1/2})$. Using standard concentration inequalities we get that with high probability $d_{F'}(e) = (1 + o(1))n^{1/2}$ for each $e \in V(F')$ and that $\Delta_2(F') = O(\log n)$. This means that

$$\delta_1(F') \geq (1 - o(1))\Delta_1(F'), \quad g(F') = \Omega(n^{1/2}/\log n) \quad \text{and} \quad g(F') = O(n^{1/2}).$$

For each $(k-1)$ -set S of vertices of H , let $U_S \subseteq V(F)$ correspond to the edges in H containing S . There are at most n^{k-1} such sets and each has size at least εn . Thus, it is easy to check that the conditions for Theorem 9.7 are satisfied. Therefore, there is a matching M in F' such that at most γn vertices in $V(F')$ are uncovered in each U_S . The matching M in $F' \subseteq F$ translates to a $C_\ell^{(k)}$ -packing \mathcal{C} in H and the latter condition implies $\Delta_{k-1}(H - \bigcup \mathcal{C}) \leq \gamma n$, as desired. \blacksquare

9.3. Extending lemma. For this section we will use the following result (see [14, Theorem 5.5]).

Theorem 9.8. *Let X_1, \dots, X_t be Bernoulli random variables (not necessarily independent) such that, for each $i \in [t]$, we have $\mathbb{P}[X_i = 1 | X_1, \dots, X_{i-1}] \leq p_i$. Let Y_1, \dots, Y_t be independent Bernoulli random variables such that $\mathbb{P}[Y_i = 1] = p_i$ for all $i \in [t]$. Let $X = \sum_{i \in [t]} X_i$ and $Y = \sum_{i \in [t]} Y_i$. Then $\mathbb{P}[X \geq k] \leq \mathbb{P}[Y \geq k]$ for all $k \in \{0, 1, \dots, t\}$.*

Let \mathcal{S} be a multiset of ordered $(k-1)$ -tuples in an n -vertex set V , possibly with repetitions. We say that \mathcal{S} is γ -sparse if the multi- $(k-1)$ -graph S formed by all the unoriented $(k-1)$ -sets from \mathcal{S} , counting repetitions, has $\Delta_j(S) \leq \gamma n^{k-j}$ for each $0 \leq j \leq k-1$. For instance, the $j=1$ case says that no vertex is in more than γn^{k-1} tuples (counting repetitions). Recall the definition of ends of a trail P and $D(P)$ in Section 6.1.

Lemma 9.9 (Extending lemma). *Let $1/n \ll \gamma \ll \mu \ll \varepsilon, 1/\ell, 1/k$. Let H be a k -graph on n vertices. Let $\mathcal{S} = \{\mathbf{a}_i, \mathbf{b}_i : i \in [t]\}$ be a multiset of ordered $(k-1)$ -tuples in $V(H)$ such that*

- (a) \mathcal{S} is γ -sparse and
- (b) for each $i \in [t]$, there are at least εn^ℓ trails P in H on $\ell + 2(k-1)$ vertices such that $D(P) = \{\mathbf{a}_i, \mathbf{b}_i\}$.

Then, there exist edge-disjoint trails P_1, \dots, P_t in H such that, for each $i \in [t]$,

- (i) P_i has $\ell + 2(k-1)$ vertices and $D(P_i) = \{\mathbf{a}_i, \mathbf{b}_i\}$,
- (ii) the vertices of P_i outside \mathbf{a}_i and \mathbf{b}_i are all distinct and
- (iii) $\Delta_{k-1}(\bigcup_{i \in [t]} P_i) \leq \mu n$.

Proof. The idea is to pick, sequentially, a trail P_i chosen uniformly at random among all the trails whose ends are \mathbf{a}_i and \mathbf{b}_i . Since \mathcal{S} is γ -sparse and there are plenty of choices for P_i in each step, we expect that in each step the random choices do not affect the codegree of the graph formed by the yet unused edges in H by much. This will ensure that, even after removing the edges used by P_1, \dots, P_{i-1} , there are still many trails P_i available for the i th step. If all goes well, then we can continue the process until the end, thus finding the required trails.

We say that a trail P is i -good if P is on $\ell + 2(k-1)$ vertices, $D(P) = \{\mathbf{a}_i, \mathbf{b}_i\}$ and the vertices of P outside \mathbf{a}_i and \mathbf{b}_i are all distinct. Let $\mathcal{P}_i(H)$ be the set of all i -good trails in H . We begin by noting that $\mathcal{P}_i(H)$ is large. Indeed, there are at most $\ell^2 n^{\ell-1} \leq \varepsilon n^\ell / 2$ trails P on $\ell + 2(k-1)$ vertices with $D(P) = \{\mathbf{a}_i, \mathbf{b}_i\}$ that is not i -good. By (b), $|\mathcal{P}_i(H)| \geq \varepsilon n^\ell / 2$. Since $\mu \ll \varepsilon$, we have

$$(9.2) \quad \text{if } G \text{ is a } k\text{-graph with } \Delta_{k-1}(G) \leq \mu n, \text{ then } |\mathcal{P}_i(H - G)| \geq \varepsilon n^\ell / 3.$$

We now describe the random process. For each $i \in [t]$, assume we have already chosen edge-disjoint $P_1, P_2, \dots, P_{i-1} \subseteq H$, and we describe the choice of P_i . Let $G_{i-1} = \bigcup_{j \in [i-1]} P_j$ correspond to the edges of H used by the previous choices of P_j , which we need to avoid when choosing P_i (note that G_0 is empty). If $\Delta_{k-1}(G_{i-1}) \leq \mu n$, then (9.2) implies that $|\mathcal{P}_i(H - G_{i-1})| \geq \varepsilon n^\ell / 3$ and we take $P_i \in \mathcal{P}_i(H - G_{i-1})$ uniformly at random. Otherwise, if $\Delta_{k-1}(G_{i-1}) > \mu n$, then let $P_i = \emptyset$.

In any case, the process outputs a collection of edge-disjoint subgraphs P_1, \dots, P_t . Our task now is to show that with positive probability, there is a choice of P_1, \dots, P_t such that $\Delta_{k-1}(G_t) \leq \mu n$. This will imply also that $P_i \in \mathcal{P}_i$, which is what we needed. Formally, for each $i \in [t]$, let \mathcal{S}_i be the event that $\Delta_{k-1}(G_i) \leq \mu n$. Thus it is enough to show $\mathbb{P}[\mathcal{S}_t] > 0$.

Fix $e \in \binom{V(H)}{k-1}$. For each $i \in [t]$, let $X_i(e)$ be the random variable that takes the value 1 precisely if e belongs to an edge of P_i , and 0 otherwise. Equivalently, $X_i(e) = 1$ if and only if $\deg_{P_i}(e) \geq 1$. Since $\Delta_{k-1}(P_i) \leq 2$ for each $i \in [t]$, we have

$$(9.3) \quad \deg_{G_i}(e) \leq 2 \sum_{j \in [i]} X_j(e).$$

For each $i \in [t]$, define

$$r_i(e) = \max\{|e \cap \mathbf{a}_i|, |e \cap \mathbf{b}_i|\} \quad \text{and} \quad p_i^*(e) = \min \left\{ 1, \frac{6\ell k}{\varepsilon n^{(k-1)-r_i(e)}} \right\},$$

where here $\mathbf{a}_i, \mathbf{b}_i$ are taken as the underlying $(k-1)$ -sets.

Claim 9.10. *For each $e \in \binom{V(H)}{k-1}$ and $i \in [t]$, $\mathbb{P}[X_i(e) = 1 | X_1(e), X_2(e), \dots, X_{i-1}(e)] \leq p_i^*(e)$.*

Proof of the claim. Fix $e \in \binom{V(H)}{k-1}$ and $i \in [t]$. Using conditional probabilities, we separate our analysis depending on whether \mathcal{S}_{i-1} holds or not. If \mathcal{S}_{i-1} fails, then $P_i = \emptyset$ and so $X_i(e) = 0$ implying that our claim holds.

Now assume that \mathcal{S}_{i-1} holds, so $\Delta_{k-1}(G_{i-1}) \leq \mu n$. By (9.2), P_i will be chosen uniformly at random from $\mathcal{P}_i(H - G_{i-1})$, which has size at least $\varepsilon n^\ell / 3$ regardless of the values of $X_1(e), \dots, X_{i-1}(e)$.

If $r_i(e) = k-1$, then $p_i^*(e) = 1$ and we are done. We may assume that $r = r_i(e) \in [k-2] \cup \{0\}$. We now estimate the number of $P \in \mathcal{P}_i(H - G_{i-1})$ with $\deg_P(e) \geq 1$. If we have $P = v_1 v_2 \dots v_{\ell+2(k-1)}$ and $\deg P(e) \geq 1$, then $j_0 = \min\{j : v_j \in e\} \in [\ell+k-1]$ and $|\{j \in [k] : v_{j_0+j} \notin e\}| = 1$. Recall that, for each $P \in \mathcal{P}_i(H - G_{i-1})$, it holds that $|V(P) \setminus \{\mathbf{a}_i, \mathbf{b}_i\}| = \ell$ and it also holds that $|e \setminus \mathbf{a}_i|, |e \setminus \mathbf{b}_i| \geq k-1-r_i(e)$. Hence, we deduce that the number of $P \in \mathcal{P}_i(H - G_{i-1})$ with $\deg_P(e) \geq 1$ is certainly at

most $(\ell + k - 1)kn^{\ell - (k-1-r_i(e))} \leq 2\ell kn^{\ell - (k-1-r_i(e))}$. Thus we have

$$\mathbb{P}[X_i(e) = 1 | X_1(e), \dots, X_{i-1}(e), \mathcal{S}_{i-1}] \leq \frac{2\ell kn^{\ell - (k-1-r_i(e))}}{|\mathcal{P}_i(H - G_{i-1})|} \leq \frac{6\ell k}{\varepsilon n^{k-1-r_i(e)}} = p_i^*(e),$$

as required. This finishes the proof of the claim. \square

Now, we use that \mathcal{S} is γ -sparse to argue that $\sum_{i \in [t]} p_i^*(e)$ is small for each $e \in \binom{V(H)}{k-1}$. Indeed, for each $0 \leq r \leq k-1$, let t_r be the number of $i \in [t]$ such that $r_i(e) = r$. Since \mathcal{S} is γ -sparse, we have $t_r \leq 2 \binom{k-1}{r} \gamma n^{k-r}$ for each $0 \leq r \leq k-1$. Recall that we are assuming the hierarchy $\gamma \ll \mu \ll \varepsilon, 1/\ell, 1/k$. Therefore, we have

$$(9.4) \quad \sum_{i \in [t]} p_i^*(e) = t_{k-1} + \sum_{0 \leq r \leq k-2} t_r \cdot \frac{6\ell k}{\varepsilon n^{k-1-r}} \leq \frac{\mu}{30} n.$$

We now claim that

$$(9.5) \quad \mathbb{P} \left[\sum_{i \in [t]} X_i(e) \geq \frac{\mu}{3} n \right] \leq \exp \left(-\frac{\mu}{3} n \right).$$

Indeed, (9.4) implies that $7 \sum_{i \in [t]} p_i^*(e) \leq \mu n/3$, so the bound follows from Theorem 9.8 combined with a Chernoff-type bound [10, Corollary 2.4].

For each $e \in \binom{V(H)}{2}$, let $X_e := \sum_{i \in [t]} X_i(e)$. Let \mathcal{E} be the event that $\max_e X_e \leq \mu n/3$. By using a union bound over all the (at most n^{k-1}) possible choices of e and using (9.5), we deduce that \mathcal{E} holds with probability at least $1 - o(1)$.

Now we can show that \mathcal{S}_t holds with positive probability. In fact, we shall prove that $\mathbb{P}[\mathcal{S}_t | \mathcal{E}] = 1$, which then will imply $\mathbb{P}[\mathcal{S}_t] \geq \mathbb{P}[\mathcal{S}_t | \mathcal{E}] \mathbb{P}[\mathcal{E}] \geq 1 - o(1)$. So assume \mathcal{E} holds, that is, $\max_e X_e \leq \mu n/3$. Note that \mathcal{S}_0 holds deterministically, and suppose that $i \in [t]$ is the minimum such that \mathcal{S}_i fails to hold. Since \mathcal{S}_{i-1} holds, using (9.3) we deduce

$$\begin{aligned} \Delta_{k-1}(G_i) &\leq 2 + \Delta_{k-1}(G_{i-1}) = 2 + \max_e \deg_{G_{i-1}}(e) \leq 2 \left(1 + \max_e \sum_{j \in [i-1]} X_j(e) \right) \\ &\leq 2 \left(1 + \max_e X_e \right) \leq 2 \left(1 + \frac{\mu}{3} n \right) \leq \mu n, \end{aligned}$$

where in the penultimate inequality we used \mathcal{E} , and in the last inequality we used $1/n \ll \mu$. Thus \mathcal{S}_i holds, a contradiction. \blacksquare

The following corollary of Lemma 9.9 allows us to find a sparse path-decomposable subgraph whose removal adjusts the degrees modulo k . This was used in proving Corollary 2.4.

Corollary 9.11. *Let $0 < 1/n \ll \mu \ll 1/\ell, 1/k, \varepsilon$ with $\ell > k \geq 3$. Let H be a k -graph on n vertices such that $\delta^{(2)}(H) \geq \varepsilon n$. Then there exists a $P_\ell^{(k)}$ -decomposable subgraph H' such that*

- (i) $|H'| \leq \ell^2 kn$,
- (ii) $\Delta_{k-1}(H') \leq \mu n$ and
- (iii) for each $x \in V(H)$, we have $\deg_{H-H'}(x) \equiv 0 \pmod k$.

Proof. We start by finding a multidigraph D on $V = V(H)$ such that $d_D^+(v) - d_D^-(v) + \deg_H(v) \equiv 0 \pmod k$ holds for each $v \in V$. This can be constructed greedily, starting from an empty digraph D . As long as there is a pair of vertices u, v and $0 < i \leq j < k$ with $d_D^+(u) - d_D^-(u) + \deg_H(u) \equiv i \pmod k$ and $d_D^+(v) - d_D^-(v) + \deg_H(v) \equiv j \pmod k$, we pick them by minimising i and maximising j , and then we add the directed edge $u \rightarrow v$ to D . Since $\sum_{x \in V(H)} \deg_H(x) = k|H| \equiv 0 \pmod k$, this process is guaranteed to end. By construction, we have $d_D^+(v), d_D^-(v) \leq k$ for each $v \in V$, and D has at most kn arcs.

Let ℓ_0 be the minimum integer divisible by ℓ such that $\ell_0 \geq k^2 - k + 2$. We clearly have the inequality $\ell_0 \leq \ell^2$. Given vertices $u, v \in V$, suppose that $T = T_{u,v} \subseteq H$ is a $P_\ell^{(k)}$ -decomposable subgraph on ℓ_0 edges such that $\deg_T(u) = k - 1$, $\deg_T(v) = 1$ and $\deg_T(w) \equiv 0 \pmod k$ for all other vertices. Suppose we can find an edge-disjoint collection \mathcal{T} of such subgraphs $T_{u,v}$, one for each edge $u \rightarrow v$ in D , with the additional condition that the union H' of those subgraphs has codegree at most μn . Then H' is easily seen to satisfy the required conditions. We now describe the construction of such a family.

Each $T_{u,v}$ will be chosen as follows. Given $uv \in E(D)$, we pick a $(k - 2)$ -tuple of vertices $\mathbf{x}(u, v) = x_2 \cdots x_{k-1} \in (V \setminus \{u, v\})^{k-2}$, uniformly at random. Then, we consider the $(k - 1)$ -tuples $\mathbf{v}_{u,v} = ux_{k-1} \cdots x_2$ and $\mathbf{w}_{u,v} = x_2 \cdots x_{k-1}v$. Note that a trail with ends $\mathbf{v}_{u,v}$ and $\mathbf{w}_{u,v}$ using ℓ_0 edges and no new repeating vertices forms a $T_{u,v}$ with the required characteristics. In particular, $T_{u,v}$ has a $P_\ell^{(k)}$ -decomposition.

Consider the multiset of ordered $(k - 1)$ -tuples $\mathcal{Q} = \bigcup_{uv \in E(D)} \{\mathbf{v}_{u,v}, \mathbf{w}_{u,v}\}$. Since the bounds $\Delta^+(D), \Delta^-(D) \leq k$ hold and $\mathbf{x}(u, v)$ was chosen at random for each directed edge $uv \in E(D)$, we can assume that \mathcal{Q} is γ -sparse. Select a new constant ρ which satisfies the hierarchy $\mu \ll \rho \ll \varepsilon$. By Lemma 6.1, for each $uv \in E(D)$, there exist $\rho n^{\ell_0 - k + 1}$ trails with ℓ_0 edges and ends $\mathbf{v}_{u,v}$ and $\mathbf{w}_{u,v}$. Then, Lemma 9.9 (with ρ in place of ε) provides us with an edge-disjoint collection of trails $\{T_{uv} : uv \in E(D)\}$, one for each $uv \in E(D)$, such that T_{uv} has ends $\mathbf{v}_{u,v}$ and $\mathbf{w}_{u,v}$, no repeated vertices save for those in the ends, and $H' = \bigcup_{uv \in E(D)} T_{uv}$ satisfies $\Delta_{k-1}(H') \leq \mu n$, which is all we needed. \blacksquare

9.4. Cover-down lemma: Proof of Lemma 9.1. For this section, we will require a few pieces of new notation. Given a k -graph H , a vertex set $U \subseteq V(H)$, a $(k - 1)$ -tuple $e \in \binom{V(H)}{k-1}$, and a set of $(k - 1)$ -tuples $G \subseteq \binom{V(H)}{k-1}$, define $N_H(e, U; G) = N(e, U) \cap G$. Moreover, define

$$\delta^{(2)}(H; U, G) = \min \left\{ N_H(e_1, U; G) \cap N_H(e_2, U; G) : e_1, e_2 \in \binom{V(H)}{k-1} \right\}.$$

Proof of Lemma 9.1. Given $k \in \mathbb{N}$ and $\alpha > 0$ take $\ell_0 \in \mathbb{N}$ larger than the one given by (\curvearrowright_{k-1}) and sufficiently large for an application of Lemma 9.5. Moreover, for $\mu > 0$ we take auxiliary variables γ, p_i and μ_i for every $i \in [k - 1]$, under the following hierarchy

$$0 < \gamma \ll \mu_1 \ll p_1 \ll \cdots \ll \mu_{k-1} \ll p_{k-1} \ll \mu_k \ll \mu, \alpha.$$

Finally take $\gamma_i = \gamma + 2p_i \ll \mu_{i+1}$ and $\alpha_i = p_i \alpha^2 / 2 - \sum_{0 \leq j \leq i-1} \mu_j \gg \mu_i$. Let $n \in \mathbb{N}$ be sufficiently large and let H be as in the statement of the lemma.

Step 1: Setting the stages. For every $0 \leq i \leq k - 1$, let $H_i = \{e \in H : |e \cap U| = i\}$ and let $R_i \subseteq H_i$ be defined by choosing edges independently at random from H_i with probability p_i . Moreover, let $R_{\geq i} = \bigcup_{i \leq j \leq k-1} R_j$. Considering (CD_2) , by standard concentration inequalities we have that with non-zero probability the following inequalities happen simultaneously: for every $0 \leq i \leq k - 1$,

$$(9.6) \quad \Delta_{k-1}(R_i) \leq 2p_i n$$

$$(9.7) \quad \delta_{k-1}^{(2)}(R_{\geq i} \cup H[U]; U, G_{i-1}) \geq \frac{p_i \alpha |U|}{2} \geq \frac{p_i \alpha^2 n}{2},$$

where $G_i = \{e \in \binom{V}{k-1} : |e \cap U| \geq i\}$ (we include the degenerate cases $G_{-1} = G_0 = \binom{V}{k-1}$). From now on for every $0 \leq i \leq k - 1$ we consider R_i to be a fixed graph with those properties.

Define $H^* = H - H[U] - R_{\geq 0}$ and observe that $\delta^{(2)}(H^*) \geq \alpha n / 2$. Hence we can apply Lemma 9.5 to find a $C_\ell^{(k)}$ -packing \mathcal{C} such that $\Delta_{k-1}(H^* - \bigcup \mathcal{C}) \leq \gamma n$. We shall find a $C_\ell^{(k)}$ -packing that covers the leftover $J = H^* - \bigcup \mathcal{C}$ and the graph $R_{\geq 0}$. We

do this in stages, covering the edges $J_i = (J \cap H_i) \cup R_i$ (and some from $R_{\geq i}$) in each stage.

Step 2: The first $k - 1$ stages. To start, let $\mathcal{C}_{-1} = J_{-1} = \emptyset$. Let $0 \leq i < k - 2$ and denote the edges which were covered in previous stages by $J_{\leq i-1} = \bigcup_{0 \leq j \leq i-1} J_j$. Suppose there is a $C_\ell^{(k)}$ -packing \mathcal{C}_{i-1} such that

$$(9.8) \quad \bigcup \mathcal{C}_{i-1} \cap H[U] = \emptyset, \quad J_{\leq i-1} \subseteq \bigcup \mathcal{C}_{i-1}, \quad \text{and} \quad \Delta_{k-1}(\bigcup \mathcal{C}_{i-1} - J_{\leq i-1}) \leq \sum_{0 \leq j \leq i} \mu_j n.$$

Note that (9.8) holds vacuously for $i = 0$. We shall prove the existence of a packing \mathcal{C}_i satisfying (9.8) for i instead of $i - 1$.

Consider $\tilde{R}_{\geq i+1} = R_{\geq i+1} - \bigcup \mathcal{C}_{i-1}$ and $\tilde{J}_i = J_i - \bigcup \mathcal{C}_{i-1}$ the remaining edges from $R_{\geq i+1}$ and J_i after deleting $\bigcup \mathcal{C}_{i-1}$. Because of (9.7) and (9.8) we have that

$$(9.9) \quad \delta^{(3)}(\tilde{R}_{\geq i+1}; U, G_i) \geq \left(\frac{p_{i+1} \alpha^2}{2} - \sum_{0 \leq j \leq i+1} \mu_j \right) n = \alpha_{i+1} n \gg \mu_{i+1} n.$$

Moreover, in view of (9.6), we obtain

$$(9.10) \quad \Delta_{k-1}(\tilde{J}_i) \leq \Delta_{k-1}(J) + \Delta_{k-1}(R_i) \leq \gamma n + 2p_i \alpha n = \gamma_i n.$$

Enumerate edges of \tilde{J}_i into e_1, \dots, e_t . For each $j \in [t]$, we oriented e_j arbitrarily and let $\{\mathbf{a}_j, \mathbf{b}_j\}$ be such that $D(e_j) = \{\mathbf{a}_j^{-1}, \mathbf{b}_j^{-1}\}$. Note that $\mathcal{S} = \{\mathbf{a}_j, \mathbf{b}_j : j \in [t]\}$ is γ_i -sparse. Moreover, (9.9) and Lemma 6.1 implies, for each $j \in [t]$, \tilde{R}_{i+1} contains at least $\alpha_{i+1} n^{\ell-k}$ trails P on $\ell + k - 2$ vertices such that $D(P) = \{\mathbf{a}_j, \mathbf{b}_j\}$. We apply Lemma 9.9 with $\alpha_{i+1}, \mu_{i+1}, \gamma_i, \ell - k, \tilde{R}_{i+1}$ in the rôles of $\alpha, \mu, \gamma, \ell, H$ to obtain edge-disjoint trails P_1, \dots, P_t in \tilde{R}_{i+1} such that, for each $j \in [t]$,

- (i) P_j has $\ell + k - 2$ vertices and $D(P_j) = \{\mathbf{a}_j, \mathbf{b}_j\}$,
- (ii) the vertices of P_j outside \mathbf{a}_j and \mathbf{b}_j are all distinct and
- (iii) $\Delta_{k-1}(\bigcup_{j \in [t]} P_j) \leq \mu_{i+1} n$.

Note that $e_i \cup P_i$ is $C_\ell^{(k)}$, so $\tilde{J}_i \cup \bigcup_{j \in [t]} P_j$ has a $C_\ell^{(k)}$ -decomposition \mathcal{C}'_i . It is easy to see that by taking $\mathcal{C}_i = \mathcal{C}_{i-1} \cup \mathcal{C}'_i$ we obtain a $C_\ell^{(k)}$ -packing satisfying (9.8) with i instead of $i - 1$.

Step 3: The last stage. For the last stage, a few changes are needed. This is because in the previous stages we used edges from H_{i+1} to extend paths in H_i , which is no longer possible at this stage. Instead, we rely on the path decompositions ensured by (\curvearrowright_{k-1}) .

As before, we define $\tilde{J}_{k-1} = J_{k-1} - \bigcup \mathcal{C}_{k-2}$. For every vertex $v \in V(H) \setminus U$, we let $F(v) = L_v \cap \tilde{J}_{k-1}$ be the link graph of v in the hypergraph \tilde{J}_{k-1} . Note that $F(v)$ is completely contained in U . We shall apply (\curvearrowright_{k-1}) to find a $P_k^{(k-1)}$ -decomposition in $F(v)$. For this, we first prove that $|F(v)| = \deg_{\tilde{J}_{k-1}}(v)$ is divisible by k . Indeed, (CD₃) says that $\deg_H(v)$ is divisible by k , and since $\tilde{J}_{k-1} = H - H[U] - \bigcup \mathcal{C} - \bigcup \mathcal{C}_{k-2}$ we have $\deg_{\tilde{J}_{k-1}}(v)$ is divisible by k as well. Moreover, because of (9.7) and (9.9) we have that

$$\delta^{(3)}(F(v)) \geq \frac{p_{k-1} \alpha^2}{2} n - \sum_{0 \leq j \leq k-1} \mu_j n \geq \alpha_{k-1} n.$$

Hence, (\curvearrowright_{k-1}) yields a $P_k^{(k-1)}$ -decomposition of $F(v)$. Notice that each path in this decomposition corresponds to a $P_{k+1}^{(k)}$ in \tilde{J}_{k-1} when we include the vertex v in every edge. Call this $P_{k+1}^{(k)}$ -packing \mathcal{P}_v and observe that paths from \mathcal{P}_v and \mathcal{P}_u are edge-disjoint for every $u \neq v$. This means $\mathcal{P} = \bigcup_{v \in V(H) \setminus U} \mathcal{P}_v$ is a $P_{k+1}^{(k)}$ -decomposition of \tilde{J}_{k-1} .

Now we continue as in the previous stages and observe that

$$\Delta_{k-1}(\tilde{J}_{k-1}) \leq \Delta_{k-1}(J) + \Delta_{k-1}(R_{k-1}) \leq \gamma n + 2p_{k-1}\alpha n = \gamma_k n,$$

which implies that $D(\mathcal{P})$ (without simplification) is γ_k -sparse. Moreover, (9.8) implies that

$$\delta^{(3)}((H - \mathcal{C}_{k-2})[U]) = \delta^{(3)}(H[U]) \geq \alpha|U|,$$

By Lemma 6.1, for any $P \in \mathcal{P}$ with $D(P) = \{\mathbf{a}^{-1}, \mathbf{b}^{-1}\}$, $(H - \mathcal{C}_{k-2})[U]$ contains at least $\alpha n^{\ell-k-1}$ trails Q on $\ell + k - 3$ vertices such that $D(Q) = \{\mathbf{a}, \mathbf{b}\}$. Finally, we apply Lemma 9.9 as in the previous stages to obtain edge-disjoint trails $\{Q_P : P \in \mathcal{P}\}$ in $(H - \mathcal{C}_{k-2})[U]$ such that, for each $P \in \mathcal{P}$,

- (i) Q_P has $\ell + k - 3$ vertices and $D(Q_P) = \{\mathbf{a}, \mathbf{b}\}$ such that $D(P) = \{\mathbf{a}^{-1}, \mathbf{b}^{-1}\}$,
- (ii) the vertices of Q_P outside $D(Q_P)$ are all distinct and
- (iii) $\Delta_{k-1}(\bigcup_{P \in \mathcal{P}} Q_P) \leq \mu_k n$.

Each $P \cup Q_P$ forms a $C_\ell^{(k)}$, so $\tilde{J}_{k-1} \cup \bigcup_{P \in \mathcal{P}} Q_P = \bigcup_{P \in \mathcal{P}} (P \cup Q_P)$ has a $C_\ell^{(k)}$ -decomposition \mathcal{C}'_{k-1} . Thus, recalling (9.8), it is easy to see that the $C_\ell^{(k)}$ -packing $\mathcal{C}^* = \mathcal{C} \cup \mathcal{C}_{k-2} \cup \mathcal{C}'_{k-1}$ satisfies the requirements of the lemma. \blacksquare

10. EULERIAN TOURS

We first show that a lower bound of (essentially) $n/2$ on the codegree of k -graphs is necessary to ensure that every edge is in some tight cycle. The bound is asymptotically tight by Lemma 6.1 (which can be used to find cycles which contain any given edge). This also provides the lower bound in Theorem 1.5.

Proposition 10.1. *For all $k \geq 3$ and $m \geq 2$, there exists a k -graph H on $n = 2mk$ vertices with $\delta(H) \geq n/2 - 2k + 1$ such that $\deg(v)$ is divisible by k for all $v \in V(H)$ and there is an edge that is not contained in any tight cycle. In particular, we have the bounds $\delta_{\text{cycle}}^{(k)}, \delta_{\text{Euler}}^{(k)} \geq 1/2$.*

Proof. Let A and B be disjoint vertex-sets each of size mk . Recall that, for $0 \leq i \leq k$, we defined $H_i = H_i^{(k)}(A, B)$ as the k -graph with vertex set $A \cup B$ such that $e \in H_i$ if and only if $|e \cap B| = i$. Consider the k -graph

$$H^* = \bigcup_{i \in (\{0\} \cup [k]) \setminus \{1, k-1\}} H_i^{(k)}(A, B),$$

and observe that $\delta(H^*) \geq n/2 - k + 1$. Note that each vertex has the same vertex-degree. By removing at most $k - 1$ perfect matchings in each of $H^*[A] = H_0(A, B)$ and $H^*[B] = H_k(A, B)$, we may assume that each vertex has vertex-degree divisible by k . Additionally, remove edges $a_1 \cdots a_k \in H^*[A]$ and $b_1 \cdots b_k \in H^*[B]$ and add two edges $a_1 \cdots a_{k-1} b_k$ and $b_1 \cdots b_{k-1} a_k$. Call the resulting graph H . Note that the bound $\delta(H) \geq n/2 - 2k + 1$ holds, and for every vertex v , $d_H(v)$ is divisible by k .

We now claim that the edge $a_1 \cdots a_{k-1} b_k$ is not contained in any tight cycle. Indeed, for $k = 3$, note that $\deg_H(a_1 b_3) = \deg_H(a_2 b_3) = 1$, so $a_1 a_2 b_3$ can only be the end of any tight path implying that $a_1 a_2 b_3$ is not contained in any tight cycle. Now assume that $k \geq 4$. Since $a_1 \cdots a_{k-1} b_k$ is the only edge in $H \cap H^1(A, B)$ (i.e. with exactly $k - 1$ vertices in A) any tight path of length containing $a_1 \cdots a_{k-1} b_k$ internally must travel from $H^0(A, B)$ to $\bigcup_{i \geq 2} H^i(A, B)$. However, there is no other edge in $H \cap H^1(A, B)$ to close this tight path into a cycle.

Since we have ensured every degree in H is divisible by k , this construction shows that $\delta_{\text{cycle}}^{(k)}, \delta_{\text{Euler}}^{(k)} \geq 1/2$. \blacksquare

We split the other inequalities in Theorem 1.5 into several lemmas.

Lemma 10.2. *For $k \geq 3$, $\delta_{\text{Euler}}^{(k)} \leq \delta_{\text{cycle}}^{(k)}$.*

Proof. Let $\ell = k^2$ and $k \geq 3$. Let $0 < 1/n \ll \gamma \ll \mu \ll \varepsilon$. Let H be a k -graph on n vertices with $\delta(H) \geq (\delta_{\text{cycle}}^{(k)} + \varepsilon)n$ such that $\deg_H(v)$ is divisible by k for all $v \in V(H)$. Note that $\delta(H) \geq (1/2 + \varepsilon)n$ by Proposition 10.1. Let $\sigma_1, \dots, \sigma_t$ be an enumeration of all ordered $(k-1)$ -tuples of $V(H)$, so $t = n!/(n-k+1)!$. For each $i \in [t]$, let $\mathbf{a}_i = \sigma_i$ and $\mathbf{b}_i = \sigma_{i+1}^{-1}$, with indices taken modulo t . Let $\mathcal{S} = \{\mathbf{a}_i, \mathbf{b}_i : i \in [t]\}$ be the multisets. Note that \mathcal{S} is γ -sparse. By Lemma 6.1, for all $i \in [t]$, H contains at least εn^ℓ trails P on $\ell + 2(k-1)$ vertices such that $D(P) = \{\mathbf{a}_i, \mathbf{b}_i\}$. Apply Lemma 9.9 to obtain edge-disjoint trails $\{P_i : i \in [t]\}$ in H such that, for each $i \in [t]$,

- (i) P_i has $\ell + 2(k-1)$ vertices and $D(P_i) = \{\mathbf{a}_i, \mathbf{b}_i\}$;
- (ii) the vertices of P_i outside $D(P_i)$ are all distinct and
- (iii) $\Delta_{k-1}(\bigcup_{i \in [t]} P_i) \leq \mu n$.

Let $\mathcal{P} = \bigcup_{i \in [t]} P_i$, and note that (after joining trails) we obtain a tour in H . Consider the k -graph $H' = H - \mathcal{P}$. Note that $\deg_{H'}(v)$ is divisible by k for all $v \in V(H')$ and $\delta(H') \geq \delta(H) - \mu n \geq (\delta_{\text{cycle}}^{(k)} + \varepsilon/2)n$. Thus there is a cycle-decomposition \mathcal{C} of H' . By attaching each cycle to the tour \mathcal{P} , we obtain an Eulerian tour in H . Hence we obtain $\delta_{\text{Euler}}^{(k)} \leq \delta_{\text{cycle}}^{(k)}$, as desired. \blacksquare

Lemma 10.3. *For $k \geq 3$, $\delta_{\text{cycle}}^{(k)} \leq \delta_{\text{Euler}}^{(k)}$.*

Proof (sketch). Let $\delta = \delta_{\text{Euler}}^{(k)}$, by Proposition 10.1 we have $\delta \geq 1/2$. Given $\varepsilon > 0$, let n be sufficiently large and let H be a k -graph on n vertices with $\delta(H) \geq (\delta + 2\varepsilon)n$ with all vertex-degree divisible by k . It is enough to show that H is decomposable into cycles.

The idea is to use the iterative absorption framework. Indeed, since $\delta \geq 1/2$, we have $\delta^{(2)}(H) \geq 4\varepsilon n$. Thus there exists ℓ large enough (depending on ε only) such that the Vortex lemma (Lemma 4.1) and the Cover-down lemma (Lemma 4.2) work in this setting. Thus it is possible to find a vortex $U_0 \supseteq U_1 \supseteq \dots \supseteq U_t$ to find a $C_\ell^{(k)}$ -packing which cover all edges except but those located in U_t . In fact, we can assume that the leftover $F \subseteq H[U_t]$ satisfies $\delta(F) \geq (\delta + \varepsilon)|U_t|$ (see the proof of Theorem 1.3 in Section 4 for detailed calculations to make these two steps work). The only missing step is the construction of an absorber for such a constant-sized leftover.

The key observation here is that since the leftover F will satisfy $\delta(F) \geq (\delta + \varepsilon)|U_t|$, we can assume that F admits an Euler tour. Since an Euler tour admits an edge-bijective homomorphism from a cycle, we can easily build a cycle-decomposable transformer between such a leftover and a cycle, and this step requires only $\delta^{(2)}(H) \geq \varepsilon n$ (this is exactly what is done in the proof of Lemma 4.3). \blacksquare

Lemma 10.4. *For $k \geq 3$, $\delta_{\text{cycle}}^{(k)} \leq \inf_{\ell > k} \{\delta_{C_\ell}^{(k)}\}$.*

Proof. Let $\delta = \inf_{\ell > k} \{\delta_{C_\ell}^{(k)}\}$. Note that $\delta \geq 1/2$ by Proposition 10.1.

Let $1/n \ll \varepsilon$ and let H be a k -graph on n vertices with $\delta(H) \geq (\delta + 3\varepsilon)n$ and every degree divisible by k . By the definition of infimum, there exists ℓ (depending on ε only) such that $\delta_{C_\ell}^{(k)} \leq \delta + \varepsilon$. Since $\delta(H) \geq (1/2 + 2\varepsilon)n$, we can use Lemma 6.1 to find a cycle C whose removal leaves a number of edges divisible by ℓ . Thus $\delta(H - C) \geq (\delta + 2\varepsilon)n \geq (\delta_{C_\ell}^{(k)} + \varepsilon)n$, and therefore $H - C$ admits a $C_\ell^{(k)}$ -decomposition. Together with C , this is a cycle decomposition of H . \blacksquare

Theorem 1.5 follows immediately from Lemma 10.2, Lemma 10.4, Proposition 10.1 and Theorem 1.3.

11. CONCLUDING REMARKS

Theorem 1.5 and Theorem 1.3 show that, for all k and sufficiently large ℓ , the inequalities $1/2 \leq \delta_{\text{Euler}}^{(k)} = \delta_{\text{cycle}}^{(k)} \leq \delta_{C_\ell}^{(k)} \leq 2/3$ are valid. For $k = 3$, the second and third authors [14] gave an example showing that $\delta_{\text{Euler}}^{(3)} \geq 2/3$ and thus in this the we actually have the equality $\delta_{\text{Euler}}^{(3)} = \delta_{\text{cycle}}^{(3)} = \delta_{C_\ell}^{(3)} = 2/3$ holds for all large ℓ . However,

we were unable to generalise the examples presented there for $k \geq 4$. Our best example (Proposition 10.1) gives us $\delta_{\text{cycle}}^{(k)} \geq 1/2$, so we propose the following question.

Question 11.1. *Does there exist $k \geq 4$ such that $\delta_{\text{cycle}}^{(k)} > 1/2$?*

We gave a new lower bound for the fractional $C_\ell^{(k)}$ -decomposition threshold $\delta_{C_\ell^{(k)}}^*$ in Proposition 2.1. Moreover, when $k/\gcd(\ell, k)$ is even or $\gcd(\ell, k) = 1$, we are able to calculate the value given by our bound in an explicit form (see Corollary 2.3). Is the construction given by Proposition 2.1 best-possible? We would like to propose the following weaker question.

Question 11.2. *Given $k \geq 2$, does there exist ℓ_0 such that, for all $\ell > \ell_0$ with $\ell \not\equiv 0 \pmod k$, $\delta_{C_\ell^{(k)}}^* \leq \frac{1}{2} + \frac{1}{2(\ell-1)}$?*

When $k = 2$, we believe that ℓ_0 should be 1, which also implies the Nash-Williams conjecture [13] on δ_{K_3} (c.f. [3, Theorem 1.4]).

REFERENCES

- [1] Noga Alon and Raphael Yuster. “On a hypergraph matching problem”. *Graphs Combin.* 21.4 (2005), 377–384. DOI: [10.1007/s00373-005-0628-x](https://doi.org/10.1007/s00373-005-0628-x).
- [2] Ben Barber, Stefan Glock, Daniela Kühn, Allan Lo, Richard Montgomery, and Deryk Osthus. “Minimalist designs”. *Random Structures Algorithms* 57.1 (2020), 47–63. DOI: [10.1002/rsa.20915](https://doi.org/10.1002/rsa.20915).
- [3] Ben Barber, Daniela Kühn, Allan Lo, and Deryk Osthus. “Edge-decompositions of graphs with high minimum degree”. *Adv. Math.* 288 (2016), 337–385. DOI: [10.1016/j.aim.2015.09.032](https://doi.org/10.1016/j.aim.2015.09.032).
- [4] F. Botler, G. O. Mota, M. T. I. Oshiro, and Y. Wakabayashi. “Decomposing highly edge-connected graphs into paths of any given length”. *J. Combin. Theory Ser. B* 122 (2017), 508–542. DOI: [10.1016/j.jctb.2016.07.010](https://doi.org/10.1016/j.jctb.2016.07.010).
- [5] Fan Chung, Persi Diaconis, and Ron Graham. “Universal cycles for combinatorial structures”. *Discrete Math.* 110.1-3 (1992), 43–59. DOI: [10.1016/0012-365X\(92\)90699-G](https://doi.org/10.1016/0012-365X(92)90699-G).
- [6] Stefan Glock, Felix Joos, Daniela Kühn, and Deryk Osthus. “Euler tours in hypergraphs”. *Combinatorica* 40.5 (2020), 679–690. DOI: [10.1007/s00493-020-4046-8](https://doi.org/10.1007/s00493-020-4046-8).
- [7] Stefan Glock, Daniela Kühn, Allan Lo, and Deryk Osthus. “The existence of designs via iterative absorption: hypergraph F -designs for arbitrary F ”. To appear in *Mem. Amer. Math. Soc.* Preprint (2016). DOI: [10.48550/ARXIV.1611.06827](https://doi.org/10.48550/ARXIV.1611.06827).
- [8] Stefan Glock, Daniela Kühn, and Deryk Osthus. “Extremal aspects of graph and hypergraph decomposition problems”. *Surveys in Combinatorics 2021*. Vol. 470. London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 2021, 235–265. DOI: <https://doi.org/10.1017/9781009036214.007>.
- [9] Jie Han, Allan Lo, and Nicolás Sanhueza-Matamala. “Covering and tiling hypergraphs with tight cycles”. *Combin. Probab. Comput.* 30.2 (2021), 288–329. DOI: [10.1017/S0963548320000449](https://doi.org/10.1017/S0963548320000449).
- [10] Svante Janson, Tomasz Łuczak, and Andrzej Ruciński. “Random graphs”. Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York, 2000, xii+333. DOI: [10.1002/9781118032718](https://doi.org/10.1002/9781118032718).
- [11] Felix Joos and Marcus Kühn. “Fractional cycle decompositions in hypergraphs”. *Random Structures Algorithms* (2021), 1–19. DOI: <https://doi.org/10.1002/rsa.21070>.
- [12] Peter Keevash. “The existence of designs II”. Preprint (2018). DOI: [10.48550/ARXIV.1802.05900](https://doi.org/10.48550/ARXIV.1802.05900).
- [13] C. St. J. A. Nash-Williams. “An unsolved problem concerning decomposition of graphs into triangles”. *Combinatorial Theory and its Applications III*. North Holland, 1970, 1179–183.
- [14] Simón Piga and Nicolás Sanhueza-Matamala. “Cycle decompositions in 3-uniform hypergraphs”. To appear in *Combinatorica*. Preprint (2021). DOI: [10.48550/ARXIV.2101.12205](https://doi.org/10.48550/ARXIV.2101.12205).
- [15] V. Rödl, M. Schacht, M. H. Siggers, and N. Tokushige. “Integer and fractional packings of hypergraphs”. *J. Combin. Theory Ser. B* 97.2 (2007), 245–268. DOI: [10.1016/j.jctb.2006.05.006](https://doi.org/10.1016/j.jctb.2006.05.006).
- [16] Amelia Taylor. “On the exact decomposition threshold for even cycles”. *J. Graph Theory* 90.3 (2019), 231–266. DOI: [10.1002/jgt.22399](https://doi.org/10.1002/jgt.22399).

(A. Lo, S. Piga) SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, EDGBASTON, BIRMINGHAM, B15 2TT, UK

Email address: s.a.lo@bham.ac.uk, s.piga@bham.ac.uk

(N. Sanhueza-Matamala) DEPARTAMENTO DE INGENIERÍA MATEMÁTICA, FACULTAD DE CIENCIAS FÍSICAS Y MATEMÁTICAS, UNIVERSIDAD DE CONCEPCIÓN, CHILE

Email address: nicolas@sanhueza.net